171. $I = \int_0^5 \frac{x^2}{x^2+(5-x)^2}\,dx$ ... (i)

$I = \int_0^5 \frac{(5-x)^2}{(5-x)^2+(x)^2}\,dx$ ... (ii) [f(x) = f(a–x)]

$2I = \int_0^5 dx = [x]_0^5$

$2I = 5$

$I = 5/2$

Ans. (d) None of these

172. $\lim_{x\to 0} \frac{(2^x-1)}{\sqrt{1+x}-1}.\frac{\sqrt{1+x}+1}{\sqrt{1+x}+1}$

$\lim_{x\to 0} \frac{(2^x-1)}{x} . \lim_{x\to 0} \left(\sqrt{1+x}+1\right)$

App lt.

$\Rightarrow \log 2 \times 2 \Rightarrow 2 \log 2$

Ans. (a) 2log 2

173. $\lim_{x\to 0} \frac{|x-1|}{x-1}$

RHL ... $\lim_{x\to 0+} \frac{(x-1)}{(x-1)} = 1$

LHL $\lim_{x\to 0-} \frac{-(x-1)}{(x-1)} = -1$

LHL $\neq$ RHL $\therefore$ f(x) does not exist

(c) Does not exist

174. $f(x) = \frac{x-|x|}{x}$

RHL $\lim_{x\to 0+} \frac{x-x}{x} = 0$

LHL $\lim_{x\to 0-} \frac{x-(-x)}{x} = \frac{2x}{x} = 2$

f (0) = 2

© The Institute of Chartered Accountants of India

$\Rightarrow$ RHL $\neq$ LHL $\neq$ f(0) $\quad\therefore$ f(x) is not continuous at x = 0

Ans. (b) No.

175. $y = \log_3 (\log_3 x)$

$$\frac{dy}{dx} = \frac{d}{dt}\log_3 t . \frac{d}{dx}(\log_3 x)$$

$$= \frac{1}{\log_3 x . \log_3} . \frac{1}{x . \log 3} \Rightarrow \frac{1}{x . \frac{\log x}{\log 3} . (\log 3)^2}$$

$$= \frac{1}{x . \log 3 . \log x}$$

Ans. (a) $= \dfrac{1}{x . \log 3 . \log x}$

176. C I if calculated annually.

$$CI_1 = P\left[\left(1+\frac{20}{100}\right)^2 - 1\right] = \frac{11P}{25}$$

CI if calculated semi annually.

$$CI_2 = P\left[\left(1+\frac{10}{100}\right)^4 - 1\right] = \frac{4641}{10000}P$$

$$\therefore \quad \frac{4641P}{10000} - \frac{11P}{25} = 482 \Rightarrow \frac{241P}{10000} = 482$$

$\therefore$ P = 20,000

Ans. (a) Rs. 20,000

177. Value of annuity (A) = $P\left[\dfrac{(1+i)^n - 1}{i}\right]$

$$= 3000\left[\frac{(1+0.09)^3 - 1}{(0.09)}\right]$$

= 9833.33

Ans. (c) Rs. 9833.33

© The Institute of Chartered Accountants of India

178. Present Value of Annuity $= \frac{10000}{(1+0.05)^{10}}$

= Rs. 7724

Ans. (a) Rs. 7724

179. $A = \frac{P}{i}\left[(1+i)^n - 1\right] \Rightarrow \frac{45000}{0.06}\left[(1+0.06)^{10} - 1\right]$

$750000\left[(1+0.06)^{10} - 1\right]$ {Solve by taking log}

A = 517500

$\therefore$ Surplus = 517500 – 5,00,000

= 17,500

Ans. (c) Rs. 17,500

180. Present value of P (rest) of the annuity.

$P = A\left[\frac{1-(1+i)^{-n}}{i}\right] \Rightarrow 2000\left[\frac{1-(1+0.1)^{-5}}{(0.10)}\right]$

$P = 20000\left[1-(1.1)^{-5}\right]$

P = 7294 which is less than the Purchase Price.

$\therefore$ leasing is preferable.

Ans. (a) leasing is preferable.

181. The sum of deviations of the given values from their Arithmetic Mean is 0.

Ans. (a) Arithmetic Mean

182. The sum of squares of the deviations of the given values from their Arithmetic Mean is minimum.

Ans. (a) Arithmetic Mean

183. Which is greatly affected by the extreme values – Arithmatic mean

Ans. (a) Arithmetic Mean

184. Which is not amenable to further algebric treatment – Mode and Median

Ans. (d) Both (b) and (c)

185. $\frac{a+b}{2} = 15 \Rightarrow a + b = 30$ … (i)

b – a = 4 – (ii) [By eq (i) & (ii)]

$2b = 34 \Rightarrow b = 17$

© The Institute of Chartered Accountants of India

$\therefore$ a = 13 lower limit = 13

Ans. (c) 13

186. Ans. (b) Refer Properties

187. Ans. (a) Refer Properties

188. Ans. (c) Refer Properties

189. Given, consumer price index in, (say), period I = 120 and consumer price index in (say), period II = 215. The wages of the worker in period I and II are given to be Rs. 1,680 and Rs. 3000 respectively. The real wages of the worker in the current period II with respect to the period I as base, are given by:

Rs. $\frac{120}{215} \times 3000 = \text{Rs}.1674.42$

Since this wage (Rs. 1674.42) is less than the wages of the worker in the period I (Viz. Rs. 1680) the workers is not better off but worse off R. 5.58 as compared to the period I.

Ans. (a)

190. Purchasing power (P.P) of a rupee in 1994 with respect to the base period 1980 is given by

$$\text{P.P. of a rupee} = \frac{100}{\text{Consumer Pr ice Index for 1994 w.r.t. base 1980}}$$

$$= \text{Rs. } \frac{100}{250} = \text{Re. } 0.40$$

Ans. (a)

191. Let B_1, B_2 and B_3 be the events of drawing a boy from the 1st, 2nd and 3rd group respectively and G_1, G_2 and G_3 be the events of drawing a girl from the 1st, 2nd, and 3rd group respectively then $P(B_1) = 1/4$, $P(B_2) = 2/4$, $P(B_3) = 3/4$ and $P(G_1) = 3/4$, $P(G_2) = 2/4$, $P(G_3) = 1/4$.

The required event of getting 1 girl and 2 boys in a random selection of 3 children can materialize in the following mutually exclusive cases.

(i) Girl from the first group and boys from the 2nd and 3rd group i.e. the event $G_1 \cap B_2 \cap B_3$

(ii) Girl from the 2nd group and boys from 1st and 3rd groups, i.e. the event $B_1 \cap G_2 \cap B_3$ happens.

(iii) Girl from the 3rd group and boys from the 1st and 2nd groups. i.e., the event $B_1 \cap B_2 \cap G_3$ happens.

Hence by the addition theorem of probability, required probability ρ is given by:

P = P(i) + P(ii) + P(iii)

© The Institute of Chartered Accountants of India

$= \quad P(G_1 \cap B_2 \cap B_3) + P(B_1 \cap G_2 \cap B_3) + P(B_1 \cap B_2 \cap G_3)$

$= \quad P(G_1)\,P(B_2)\,P(B_3) + P(B_1)\,P(G_2)\,P(B_3) + P(B_1)\,P(B_2)\,P(G_3)$

$= \quad \frac{3}{4}\times\frac{2}{4}\times\frac{3}{4}+\frac{1}{4}\times\frac{2}{4}\times\frac{3}{4}+\frac{1}{4}\times\frac{2}{4}\times\frac{1}{4}$

$= \quad \frac{18+6+2}{64}=\frac{26}{64}=\frac{13}{32}$

192. Let P(A) = x

$\therefore$ P(B) = $\frac{3}{2}$ P(A) = $\frac{3}{2}$x

and P(C) = $\frac{1}{2}$ P(B) = $\frac{1}{2}$ $\left(\frac{3}{2}x\right)$ = $\frac{3}{4}$x

The events A, B and C are exhaustive

$\therefore$ P (A or B or C) = 1

$\Rightarrow$ P(A) + P(B) + P(C) = 1 ($\because$ A, B, C are mutually exclusive)

$x+\frac{3}{2}x+\frac{3}{4}x=1$

$x\left[\frac{4+6+3}{4}\right]=1$

$\therefore \quad x = \frac{4}{13}$

$\therefore \quad P(A) = \frac{4}{13}$

Ans. (b)

193. There are 3+4+2+1 = 10 members in all and a Committee of 4 out of them can be formed in $10C_4$ ways. Hence exhaustive number of Cases is:

$10C_4 = \frac{10\times9\times8\times7}{4!}=210$

The probability 'p' that the Committee Consists of the doctor and at least one economist is given by

p = P [One doctor, One economist, 2 others]

+ P [One doctor, Two economist, 1 others]

+ P[One doctor, Three economist]

© The Institute of Chartered Accountants of India

$$p = \frac{1C_1 \times 3C_1 \times 6C_2}{10C_4} + \frac{1C_1 \times 3C_2 \times 6C_1}{10C_4} + \frac{1C_1 \times 3C_3}{10C_4}$$

$$= \frac{1}{210}\left[\left(1 \times 3 \times \frac{6 \times 5}{2}\right) + (1 \times 3 \times 6) + (1 \times 1)\right]$$

$$= \frac{1}{210}[45 + 18 + 1]$$

$$= \frac{64}{210} = \frac{32}{105} = 0.3048$$

Ans. (a)

194. Let A = event that the company executive travel by plane.

$\therefore$ P(A) = 2/3

Let B = event that the Company executive travel by train.

$\therefore$ P(B) = 1/5

Now the events A and B are mutually exclusive, because he cannot travel by plane and train at the same time.

$\therefore$ The prob. of his traveling by plane or train

= P (A or B) Ans. (a)

= P(A) + P(B)

= 2/3 + 1/5

$$= \frac{13}{15}$$

Ans. (b)

195. Let A and B denote the events that the contractor will get a 'plumbing' Contract and 'Electric' Contract respectively. Then we are given:

$P(A) = 2/3, P(\overline{B}) = 5/9$

$\therefore P(B) = 1 - P(\overline{B}) = 4/9$

and $P(A \cup B)$ = Probability that Contractor gets at least one contract.

= 4/5

$\Rightarrow P(A) + P(B) - P(A \cap B) = 4/5$

$2/3 + 4/9 - P(A \cap B) = 4/5$

$\Rightarrow P(A \cap B) = 2/3 + 4/9 - 4/5$

© The Institute of Chartered Accountants of India

$= \quad \frac{30+20-36}{45}$

$= \quad \frac{14}{45}$

Hence, the probability that the Contractor will get both the contracts in 14/45

Ans. (a)

196. Standard deviation = σ

$P(x > 60) = 0.05$

$\Rightarrow$ 1–P ($x \leq 60$) = 0.05

$\therefore \quad P(x \leq 60) = 0.95$

$\therefore \quad P\left[\frac{x-60}{\sigma} \leq \frac{60-50}{\sigma}\right] = 0.95$

$\therefore \quad P\left(\leq \frac{10}{\sigma}\right) = 0.95 \Rightarrow \varphi\left(\frac{10}{\sigma}\right) = \varphi(1.64)$

$\Rightarrow \quad \sigma = \left(\frac{10}{1.64}\right) = 6.7 \Rightarrow$ S.D. = 6.7

Ans. (a) 6.7

197. $P\left(Z \geq \frac{x-10}{20}\right) = 0.10$

$\therefore \quad \frac{100-x}{20} = 1.28$

$100 - x = 25.6$

$x = 74.40$

Ans. (c) 74.40

198. $P\left(x \leq \frac{x-100}{20}\right) = 0.10$

$\therefore \quad \frac{x-100}{20} = 1.28$

$x = 25.6 + 100 = 125.6$

Ans. (b) 125.6

200. P = P(x>70)

$= \quad 1 - P(x \leq 70)$

© The Institute of Chartered Accountants of India

$= \quad 1 - P\left[\frac{x-65}{25} \le \frac{70-65}{5}\right]$

$= \quad 1 - P\,(z \le 1)$

$= \quad 1\,0 - .041$

$P = 0.06$

Ans. (c) 0.06

Model Test Paper – BOS/CPT – 4

151. $P = \frac{100\,A}{100 + RT} \Rightarrow \frac{100 \times 21315}{100 + 0.045 \times \frac{4}{12}}$

$P = 21000$

Ans. (a) Rs. 21000

152. $I_1 = 500 \times \frac{8}{100} \times 1 = 40$

$I_2 = 1000 \times \frac{8}{100} \times \frac{3}{4} = 60$

$I_3 = 1000 \times \frac{8}{100} \times \frac{1}{2} = 40$

$\therefore$ Total Amount = 500 + 1000 + 1000+40 + 60 + 60

= 2640

Ans. Rs. 2640

153. $2000 = 1200\left(1 + \frac{5}{4 \times 100}\right)^{4n}$

$5 = 3\left(\frac{81}{80}\right)^{4n}$ After taking log both side and solved.

$\log 5 = \log 3 + 4n\,[\log 81 - \log 80]$

n = 10 years 3 months

Ans. (a) 10 years 3 months

© The Institute of Chartered Accountants of India

154. $26500 = 20000 \left(1+\frac{r}{100}\right)^4$

After taking log and solving it

r = 7.5%

Ans. (c) 7.5%

155. $CI = 7000 \left[\left(1+\frac{7}{100}\right)\left(1+\frac{8}{100}\right)\left(1+\frac{85}{100}\right)-1\right]$

CI = 1776

157. $I_1 = P \times \frac{3}{100} \times 2 = \frac{6P}{100}$

$I_2 = P \times \frac{8}{100} \times 3 = \frac{24P}{100}$

$I_3 = P \times \frac{10}{100} \times 1 = \frac{10P}{100}$

Total interest = 1520

$\therefore \frac{40P}{100} = 1520 \Rightarrow P = \frac{100 \times 1520}{40}$

P = 3800

Ans. Rs. 3800 (a)

158. $A = 7500 \left[(1+i)^n\right]$ [I = 0.01 n = 2]

$= 7500 [1+0.01)^2 \Rightarrow 7500 \times (1.01)^2$

A = 7650.75

Ans. (a) Rs. 7650.75

159. $512.50 = P \left[(1+0.05)^2 - 1\right]$

$512.50 = P \times 0.1025$

$\therefore$ P = 5000

Ans. (b) Rs. 5000

160. $1331 = 1000 \left[1+\frac{r}{100}\right]^3$

$\left(\frac{11}{10}\right)^3 = \left(1+\frac{r}{100}\right)^3 \Rightarrow 1.1 = 1+ \frac{r}{100}$

© The Institute of Chartered Accountants of India

$\therefore$ 0.1 = r / 100 $\Rightarrow$ r = 10%

Ans. (a) 10%

161. Range = L – S

L … S = 20 $\rightarrow$ (i)

If each item is increased by 15

Range = (L + 15) – (S + 15)

Range = L – S = 20 [from eg (i)]

Ans. (a) 20

162. Range = L – S

$\therefore$ L – S 20

If each item is divided by – 2

$$\text{Range} = \frac{L}{-2} - \frac{S}{-2} = \frac{-1}{2}(L-S)$$

$$\text{Range} = \frac{-1}{2} \times 20 = -10 \text{ [(–) sign ignored]}$$

Range = 10

(because it is difference between largest and smallest data)

Ans. (b) 10

164. In grouped frequency distribution, if the class interval is unequal then quartile deviation is more appropriate.

Ans. (a) Q.D.

165. $SD = \sqrt{\frac{\sum d^2}{n}} \Rightarrow (4)^2 = \frac{\sum d^2}{10} \Rightarrow \sum d^2 = 160$

If each item divided by – 2

Corrected $\sum (d')^2 = \frac{160}{(-2)^2} = 40$

$\therefore$ Corrected S.D. $= \sqrt{\frac{\sum (d')^2}{n}} = \sqrt{\frac{40}{10}} = 2$

S.D. = 2

Ans. (a) 2

© The Institute of Chartered Accountants of India

166. $\bar{x} = \dfrac{(5+10+15 \ldots\ldots\ldots\ldots +125}{25}$

$$\frac{\frac{25}{2}[2\times 5+(25-1)5]}{25} = \frac{130}{2} = 65$$

Average = 65

Ans. (a) 65

167. a, b, c, d, e are five add integers.

$$\text{average} = \frac{a+b+c+d+e}{5} = \frac{a+(a+2)+(a+4)+(a+6)+(a+8)}{5}$$

$\Rightarrow (a + 4)$

Ans. (d) a+4

168. $\text{Av} = \dfrac{180+258+x}{3}$

$$230 = \frac{438+x}{3} \Rightarrow 690-438 = x$$

$\therefore$ x = 252 he should score 252 runs.

Ans. (d) None of these

169. $100 = \dfrac{(x+2)60+x.120+(x-2)180}{(x+2)+x+x-2}$

$300x = 360x - 240$

$\therefore$ $60x = 240$

$x = 4$

Ans. (a) 4

170. $16 = \dfrac{\sum x}{25} \Rightarrow \sum x = 400$

$15 = \dfrac{\sum x^1}{24} \Rightarrow \sum x^1 = 360$ $\therefore$ Age of Teacher $= 400 - 360 = 40$

Ans. 40 Years

171. Ans. (b) Refer Properties

172. Ans. (b) Refer Properties

173. Given two regression lines are

© The Institute of Chartered Accountants of India

$3x + 2y = 26 \rightarrow (1)$

$6x + y = 31 \rightarrow (2)$

Since the two lines of regression intersect at the point $(\bar{x}, \bar{y})$, replacing $\bar{x}$ and $\bar{y}$ by and respectively in the given regression equation, we get.

(1) $\Rightarrow$ $2\ \bar{y} = 26 - 3\ \bar{x}$

$$\bar{y} = 13 - \frac{3}{2}\ \bar{x} \rightarrow (3)$$

(2) $\Rightarrow$ $6\ \bar{x} + 13 - \frac{3}{2}\ \bar{x} = 31$

$$\frac{12\bar{x} + 26 - 3\bar{x}}{2} = 31$$

$$9\bar{x} + 26 = 62$$

$$9\bar{x} = 62 - 26$$

$= 36$

$\bar{x} = 4$

$\therefore$ (3) $\Rightarrow$ $\bar{y} = 13 - \frac{3}{2}$ (4)

$= 13 - 6 = 7$

$\therefore$ $\bar{x} = 4,\ \bar{y} = 7$

Ans. (a)

174. Let us assume that $3x + 24 = 26 \rightarrow (1)$ represent the regression line of y on x and

$6x + y = 31 \rightarrow (2)$ represent the regression line of x on y.

(1) $\Rightarrow$ $2y = 26 - 3x$

$$y = 13 - \frac{3}{2} - x$$

$$\therefore \quad byx = -\frac{3}{2}$$

(2) $\Rightarrow$ $6x = 31 - y$

$$x = \frac{31}{6} - \frac{1}{6}\ y$$

$$\therefore \quad bxy = -\frac{1}{6}$$

© The Institute of Chartered Accountants of India

$\therefore \quad r^2 = byx \times bxy = \left(-\frac{3}{2}\right)\left(-\frac{1}{6}\right) = \frac{1}{4}$

$r = \sqrt{\frac{1}{4}} = \pm\frac{1}{2} = \pm 0.5$

Ans. (b)

We take the sign of n as negative since both the regression coefficients are negative).

175. Ans. (a) Refer Properties

176. Let E_1, E_2, E_3 denote the events that the probability is solved by X,Y and Z respectively.

Then we have

$P(E_1) = 1/3 \Rightarrow P(\overline{E}_1) = 1 - P(E_1) = 2/3$

$P(E_2) = 1/4 \Rightarrow P(\overline{E}_2) = 1 - P(E_2) = 3/4$

$P(E_3) = 1/5 \Rightarrow P(\overline{E}_3) = 1 - P(E_3) = 4/5$

Problem will be solved if at least one of the three is able to solve it. Hence, the required probability that the problem will be solved is given by $P(E_1 \cup E_2 \cup E_3)$

$= \quad 1 - P(\overline{E}_1 \cap \overline{E}_2 \cap \overline{E}_3)$

$= \quad 1 - \left[P(\overline{E}_1) \cdot P(\overline{E}_2) \cdot P(\overline{E}_3)\right]$

$= \quad 1 - 2/3 \times 3/4 \times 4/5$ [Since E_1, E_2, E_3 are independent]

$= \quad 1 - 2/5 = 3/5$

Ans. (c)

177. Given $P(A) = \frac{1}{2}$

$P(B) = \frac{1}{3}$

$P(A \cap B) = \frac{1}{4}$

$\therefore \quad P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{4}}{\frac{1}{3}} = \frac{1}{4} \times \frac{3}{1} = \frac{3}{4}$

Ans. (a)

© The Institute of Chartered Accountants of India

ANSWERS

178. Given $P(A) = \frac{1}{2}$, $P(B) = \frac{1}{3}$ and $P(A \cap B) = \frac{1}{4}$

$\therefore P(\bar{A} \cap B) = P(B) - P(A \cap B)$

$= \frac{1}{3} - \frac{1}{4} = \frac{4-3}{12} = \frac{1}{12}$

Ans. (c)

179. Given $P(A) \frac{1}{2}$, $P(B) = \frac{1}{3}$, $P(A \cap B) = \frac{1}{4}$

$P(\bar{A} \cap \bar{B}) = 1 - P(A \cup B)$

$= 1 - \{P(A) + P(B) - P(A \cap B)\}$

$= 1 - P(A) - P(B) + P(A \cap B)$

$= 1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4}$

$= \frac{12-6-4+3}{12} = \frac{5}{12}$

Ans. (a)

180. $P(A) = ½$, $P(B) = \frac{1}{3}$, $P(A \cap B) = \frac{1}{4}$

$= P(\bar{A} \cup \bar{B}) = P(\overline{A \cap B})$

$= 1 - P(A \cap B)$

$= 1 - ¼$

$= \frac{4-1}{4} = \frac{3}{4}$

Ans. (b)

181. Given $x_1 = 1, x_2 = 2, x_3 = 3$

$P(x_1) = \frac{1}{2}, P(x_2) = \frac{1}{3}, P(x_3) = 1/6$

$\therefore E(x) = x_1 P(x_1) + x_2 P(x_2) + x_3 P(x_3)$

$= \left(1 \times \frac{1}{2}\right) + \left(2 \times \frac{1}{3}\right) + \left(3 \times \frac{1}{6}\right) = \frac{1}{2} + \frac{2}{3} + \frac{1}{2}$

© The Institute of Chartered Accountants of India

$= \quad \frac{3+4+3}{6} = \frac{10}{6} = \frac{5}{3} = 1.666....$

$= \quad 1.67$

Ans. (c)

182. Given $x_1 = 1, x_2 = 2, x_3 = 3$

$P(x_1) = \frac{1}{2}, P(x_2) = \frac{1}{3}, P(x_3) = \frac{1}{6}$

$\therefore \quad V(x) = E(x^2) \ldots [E(x)]^2$

$E(x) = x_1 P(x_1) + x_2 P(x_2) + x_3 P(x_3)$

$= \quad \left(1 \times \frac{1}{2}\right) + \left(2 \times \frac{1}{3}\right) + \left(3 \times \frac{1}{6}\right)$

$E(x) = \frac{1}{2} + \frac{2}{3} + \frac{1}{2}$

$= \quad \frac{3+4+3}{6} = \frac{10}{6} = \frac{5}{3}$

$E(x^2) = x_1^2 P(x_1) + x_2^2 P(x_2) + x_3^2 P(x_3)$

$= \quad \left(1 \times \frac{1}{2}\right) + \left(4 \times \frac{1}{3}\right) + \left(9 \times \frac{1}{6}\right)$

$= \quad \frac{1}{2} + \frac{4}{3} + \frac{3}{2}$

$= \quad \frac{3+8+9}{6} = \frac{20}{6} = \frac{10}{3}$

$\therefore \quad V(x) = \frac{10}{3} - \left(\frac{5}{3}\right)^2$

$= \quad \frac{10}{3} - \frac{25}{9}$

$V(x) = \frac{30-25}{9} = \frac{5}{9} = .5556$

Ans. (a)

183. Let x denote the number of defective lamps.

X can assume the values 0, 1, 2, 3

© The Institute of Chartered Accountants of India

p(X=0) p:probably having 0 bad orange out of 4 bad orange and 3 good orange out of 8 good orange.

$$P(x=0) = \frac{4C_0 \times 8C_3}{12C_3} = \frac{56}{55}$$

$$P(x=1) = \frac{4C_1 \times 8C_2}{12C_3} = \frac{28}{55}$$

$$P(x=2) = \frac{4C_2 \times 8C_2}{12C_3} = \frac{12}{55}$$

P(x=3) = Probability having 3 bad orange out of 4 bad orange and 0 good orange out of 8 good orange.

$$= \frac{4C_3 \times 8C_0}{12C_3} = \frac{1}{55}$$

Probability that at least one orange out of three oranges is good = 1 – P (x=3)

$$= 1 - 1/55$$

$$= \frac{55-1}{55} = \frac{54}{55}$$

Ans. (a)

184. Given P(A) = 0.5, P (AB) $\leq$ 0.3

By Addition thereon,

P(A or B) = P(A) + P(B) – P (AB)

$\therefore$ P(A) + P(B) – P(AB) $\leq$ 1 [$\therefore$ P(A or B) $\leq$ 1]

$\therefore$ P(B) $\leq$ 1 – P(A) + P (AB)

$\leq$ 1 – 0.5 + 0.3

P(B) $\leq$ 0.8

Ans. (a)

185. Let the given events be A, B and P(A) = 2/3 P(B)

Let P(B) = x

$\therefore$ P(A) = 2/3 x

The events A and B are exhaustive

$\therefore$ P(A or B) = 1

P(A) + P(B) = 1

$\Rightarrow$ 2/3 x + x = 1

© The Institute of Chartered Accountants of India

5/3 x = 1

x = 3/5

∴ P(B) = 3/5

P(A) = 2/3 x 3/5 = 2/5

P(B) = 3/5 $\Rightarrow$ odds in favour of B are

3 : 5 − 3 = 3: 2

Ans. (b)

186. Given person variates with parameter = 1

i.e. $\lambda = 1$

By the poison distribution

$$p(x) = \frac{e^{-\lambda}.\lambda^x}{x!},\ x > 0$$

∴ The required probability

$$P(3 < x < 5) = P(x = 4)$$

$$= \frac{e^{-\lambda}.\lambda^4}{4!}$$

$$= \frac{e^{-1}(1)^4}{4!}$$

$$= \frac{0.36783 \times 1}{24}$$

$$P(3 < x < 5) = 0.015326$$

Ans. (a)

187. Given p = 2% = 2/100 = .02

n = 200

∴ $\lambda = np = 200 \times .02 = 4$

The probability of at least 5 defective means.

$$P(x \geq 5) = 1 - P(x < 5)$$

$$= 1 - \{P(x=0) + P(x=1) + P(x=2) + p(x=3) + P(x=4)\}$$

$$= 1 - \left\{\frac{e^{-4}(4)^0}{0!} + \frac{e^{-4}(4)^1}{1!} + \frac{e^{-4}(4)^2}{2!} + \frac{e^{-4}(4)^3}{3!} + \frac{e^{-4}(4)^4}{4!}\right\}$$

© The Institute of Chartered Accountants of India

$$= \quad 1-e^{-4}\left\{1+4+\frac{4^2}{2!}+\frac{4^3}{3!}+\frac{4^4}{4!}\right\}$$

$$= \quad 1-e^{-4}\left\{1+4+\frac{16}{2}+\frac{64}{6}+\frac{256}{24}\right\}$$

$$= \quad 1-(0.183)\ \{5+8+10.6667+10.667\}$$

$$= \quad 1-(0.83)\ (34.3334)$$

$$= \quad 1\ \ 0.6283$$

$P\ (x \geq 5) = 0.3717$

Ans. (a)

188. After a man is dealt 4 spade cards from an ordinary pack of 52 cards, there are 52 − 4 = 48 cards left in the pack, out of which 9 are spade cards and 39 are no spade cards. Now, 3 more cards can be dealt to the same man out of the 48 cards in ${}^{48}C_3$ ways, which determines the exhaustive number of ways.

If none of these 3 additional cards is a spade cards, then the 3 additional cards must be drawn out of the 39 non-spade cards, which can be done in ${}^{39}C_3$ ways.

The probability that none of the three additional cards dealt to the man is a spade card = $\frac{{}^{39}C_3}{{}^{48}C_3}$

Hence, the required probability, 'P' that at least one of the additional cards is a spade cards is given by:

$$p = \quad 1-\frac{{}^{39}C_3}{{}^{48}C_3}$$

$$= \quad 1-\frac{39\times38\times37}{3!}\times\frac{3!}{48\times47\times46}$$

$$= \quad 1-\frac{13\times19\times37}{16\times47\times23}$$

$$= \quad 1-\frac{9136}{17296}$$

$$= \quad 10.5282$$

$p = 0.4718$

Ans. (c)

190. Given P (x = 1) = P (x = 2)

Given x is a poison variable.

© The Institute of Chartered Accountants of India

$$\therefore \frac{e^{-\lambda}(\lambda)^{(1)}}{1!} = \frac{e^{-\lambda}(\lambda)^{(2)}}{2!}$$

$$\lambda = \frac{\lambda^2}{2}$$

$\lambda = 2$ = variance

Ans. (b)

191. $P = \frac{65}{500}$ $Q = 1 - P = 1 - \frac{65}{500}$

$Q = \frac{435}{500}$

$n = 500$

SE of Proportion of defectives = $\sqrt{\frac{P\,Q}{n}}$

$= \sqrt{\frac{65}{500} \times \frac{435}{500} \times \frac{1}{500}}$

SE = 0.015

Ans. (a) 0.015

192. Standard Error of Mean (SE) = $\frac{\sigma}{\sqrt{n}} = \frac{0.75}{\sqrt{100}}$

SE = 0.075

95% Confidence Limit for population mean are given by : $x \pm 1.96$ SE

$= 5.6 \pm 1.96 \times 0.75$

$= 5.6 \pm 0.147$

The Confidence level are 5.453 and 5.747

Ans. (a) 5.453 and 5.747

193. Standard Error (SE) = $\frac{\sigma}{\sqrt{n}} = \frac{4}{\sqrt{128}} = 0.353$

96% confidence limit for population mean are

$\Rightarrow \bar{x} + 2.05 \times SE$

$= 28 \pm 2.05 \times 0.353 \Rightarrow 28 \pm 0.72$

The confidence level are 27.272 and 28.728

Ans. (b) 27.272 and 28.728

© The Institute of Chartered Accountants of India

ANSWERS

194. $P = \frac{65}{500}$, $Q = 1 - P = 1 - \frac{65}{500} \Rightarrow Q = \frac{435}{500}$

SE of proportion of defectives = $\sqrt{\frac{PQ}{n}}$

$= \sqrt{\frac{65}{500} \times \frac{435}{500} \times \frac{1}{500}}$

SE = 0.015

Confidence limits for the population are

$= P \pm 3 \times SE$

$= \frac{65}{500} \pm 3 \times 0.015 \Rightarrow 0.13 \pm 0.045$

Levels are 0.085 and 0.175

or Levels are 8.5% and 17.5%

Ans. (a) 8.5% and 17.5

196. Variance = 4

$\sigma = \sqrt{4} = \pm 2$

Statement is true

Ans. (a) True

197. n = 10 P = 0.3

$\therefore Q = (1 - P) = 1 - 0.3 = 0.7$

$\therefore \sigma = \sqrt{npq}$

$\therefore$ Variance = npq = $10 \times 0.3 \times 0.7$

Variance = 2.1

Ans. (a) 2.1

198. When the cost of living increases, the standard of living improves.

Ans. (b) false

199. The 95% confidence limit for the sample mean $(\bar{x})$ is $\bar{x} \pm 1.96 \left(\frac{\sigma}{\sqrt{n}}\right)$ which is not given

Ans. (b) False

© The Institute of Chartered Accountants of India

200. Mean and variance never be equal

$\therefore$ Statement is false

Ans. (b) False

Model Test Paper – BOS/CPT – 5

151. Let the fraction be x/y. Then according to the given condition of the problem,

$$\frac{3x}{y-3}=\frac{18}{11}$$

$33x = 18y - 54$

$33x - 18y+54 = 0$

$11x - 6y+18 = 0$ ..... (i)

and $\frac{x+8}{2y}=\frac{2}{5}$

$\Rightarrow$ $5x + 40 = 4y$

$5x - 4y+40 = 0$ .....(ii)

(i) $\times 2 \Rightarrow 22x - 12y+36 = 0$ .....(iii)

(ii) $\times 3 \Rightarrow 15x - 12y+120 = 0$ ....(iv)

(iii) $-$ (iv),we get

$7x - 84 = 0$

$7x = 84$

$x = 84/7 = 12$

(i) $\Rightarrow$ $(11)(12) - 6y + 18 = 0$

$132 - 6y + 18 = 0$

$6y = 150$

$y = 150/6 = 25$

Hence, the required fraction is 12/25

$\therefore$ Ans. (c)

152. Let the two numbers are x and y

Given x + y = 150 % of y

© The Institute of Chartered Accountants of India

$= \frac{150}{100} \times y$

$x + y = 1.5y$

$x = 0.5\ y$

$x = \frac{1}{2}\ y$

Ans. (a)

153. Let three consecutive even numbers are x, x + 2, x + 4.

Given condition is $x + (x+2) + (x+4) = 60 \times \frac{3}{4} - 15$

$3x + 6 = 30$

$3x = 24$

$x = \frac{24}{3} = 8$

$\therefore$ The middle number = x + 2 = 8 + 2 = 10

Ans. (b)

154. Suppose my present age is x years and my sons present age is y years

Five years ago

my age = $(x - 5)$ years

my son's age = $(y - 5)$ years

According to the first condition of the problem,

$x - 5 = 3(y - 5)$

$x - 5 = 3y - 15$

$\Rightarrow x \quad 3y = 15 + 5$

$\Rightarrow x - 3y = -10$ .....(i)

Ten years later

my age = (x+10) years

my son's age = (y+10) years

According to the second condition of the problem,

$x+10 = 2(y+10)$

$x+10 = 2y+20$

$x-2y = 20-10$

© The Institute of Chartered Accountants of India

$x - 2y = 10$ .....(i i)

(i) − (ii) $\Rightarrow$ $y = 20$

(i) $\Rightarrow$ $x - 60 = -10$

$x = 60 - 10 = 50$

Hence, my presence age = 50 years

and my son's present age = 20 years

$\therefore$ Ans. (a)

155. The compound ratio of 4:3, 9:13, 26:5 and 2:15 is

$$= \frac{4 \times 9 \times 26 \times 2}{3 \times 13 \times 5 \times 15}$$

$$= \frac{16}{25}$$

Ans. (b)

156. We know $nPr = \frac{n!}{(n-4)!}$

$$\therefore \quad 56_{P_{r+6}} = \frac{56!}{\{56-(r+6)\}!}$$

$$= \frac{56!}{(50-r)!}$$

$$54P_{r+3} = \frac{54!}{\{54-(r+3)\}!}$$

$$= \frac{54!}{(51-r)!}$$

Thus, $\frac{56\,P_{r+3}}{54\,P_{r+3}} = \frac{56!}{(50-r)!} \times \frac{(51-r)!}{54!}$

$$= \frac{56 \times 55 \times 54!}{(50-r)!} \times \frac{(51-r)(50-r)!}{54!}$$

$$= \frac{56 \times 55 \times (51-r)}{1}$$

But we are given the ratio as 30800 : 1

$$\therefore \quad \frac{56 \times 55 \times (51-r)}{1} = \frac{30800}{1}$$

© The Institute of Chartered Accountants of India

(or) $(51 - r)! = \frac{30800}{56 \times 55} = 10 \ \therefore \ r = 41$

Ans. (b)

157. He can arrange his schedule in

$8P6 = 8 \times 7 \times 6 \times 5 \times 4 \times 3$

= 20160 ways.

Ans. (b)

158. The two Indians can stand together in ${}^2P_2 = 2! = 2$ ways.

So is the case with the two Americans and the two Russians.

Now these 3 groups of 2 each can stand in a row in ${}^3P_3 = 3 \times 2 = 6$ ways. Hence by the generalized fundamental principle, the total num ber of ways in which they can stand for a photograph under given conditions is

$6 \times 2 \times 2 \times 2 = 48$

Ans. (c)

159. This is the number of combination of 52 cards taken five at a time.

Now applying the formula.

$52C_5 = \frac{52!}{5!(52-5)!}$

$= \frac{52!}{5!47!}$

$= \frac{52 \times 51 \times 50 \times 49 \times 48 \times 47!}{5 \times 4 \times 3 \times 2 \times 1 \times 47!}$

= 2598960

Ans. (a)

160. Let the unit's digit of the number be x and the ten's digit by y. Then

$x + y = 9 \ \rightarrow (1)$

and the number = 10y + x

Reversing the order of digits of the given number,

Unit's digits becomes y

and ten's digits becomes x

$\therefore$ Now number = 10x + y

According to the given condition of the problem,

© The Institute of Chartered Accountants of India

$(10x + y) - (x+10y) = 27$

$10x + y - x - 10y = 27$

$9x - 9y = 27$

$x - y = 3 \rightarrow (2)$

Adding (1) and (2), 23 get

$2 \times = 12$

$x = \frac{12}{2} = 6$

$(1) \Rightarrow 6 + y = 9$

$y = 9 - 6 = 3$

$\therefore$ The given number is 36

Ans. (b)

161. $\underset{x\to 0}{Lt} \frac{9^x - 3^x}{4^x - 2^x} \Rightarrow \underset{x\to 0}{Lt} \left[\frac{\frac{(9^x-1)-(3^x-1)}{x}}{\frac{(4^x-1)-(2^x-1)}{x}}\right]$

$\underset{x\to 0}{Lt} \left[\frac{\left(\frac{9^x-1}{x}\right)-\left(\frac{3^x-1}{x}\right)}{\left(\frac{4^x-1}{x}\right)-\left(\frac{2^x-1}{x}\right)}\right] \Rightarrow \frac{\log 9 - \log 3}{\log 4 - \log 2}$

$\Rightarrow \frac{2\log 3 - \log 3}{2\log 2 - \log 2} = \frac{\log 3}{\log 2}$

Ans. (a) $\frac{\log 3}{\log 2}$

162. $\underset{x\to 0}{Lt} \frac{(5^x-1)^2}{\log(1+x)} = \frac{5^{2x} - 2.5^x + 1}{\log(1+x)}$

$\Rightarrow \underset{x\to 0}{Lt} \frac{(25^x - 2.5^x + 1)}{\log(1+x)} \Rightarrow \underset{x\to 0}{Lt} \frac{\left(\frac{25^x-1}{x}\right) - 2\left(\frac{5^x-1}{x}\right)}{\frac{\log(1+x)}{x}}$

App Lt

© The Institute of Chartered Accountants of India

$\Rightarrow \frac{\log 25 - 2.\log 5}{1} = \frac{\log 25 - \log 25}{1} = 0$

Ans. (d) None of these

163. $\underset{x\to 1}{Lt}\ f(x) \Rightarrow \underset{x\to 1}{Lt}\ (x+1)$

App Lt

$\Rightarrow 1 + 1 = 2$

Ans. (a) 2

164. LHL $\underset{x\to 2}{Lt}\ (x - 1)$

App Lt

$\Rightarrow 2 - 1 = 1 \quad \therefore$ LHL = 1

RHL $\underset{x\to 2}{Lt}\ (2x - 3)$

App. Lt

LHL $2.2 - 3 = 1$

$f(2) = 2.2 - 3 = 1 \qquad \therefore$ LHL = RHL = f(2)

$\therefore$ f(x) Continuous at x = 2

Ans. (a) Continuous x = 2

165. $f(x) = \frac{3x^2 + 2x + 7}{x^2 - 3x + 2} = \frac{3x^2 + 2x + 7}{(x-2)(x-1)}$

To be continuous $(x - 2) \neq 0$ & $(x - 1) \neq 0$

$\therefore x \neq 2$ & $x \neq 1$

$\therefore$ Points of discontinuity = 1, 2

Ans. (a) 1, 2

166. Let $z = \log x$

$dz = \frac{1}{x}dx$

$dx = x\,dz$

$\therefore I = \int \frac{1}{x \log x}\,dx = \int \frac{1}{x.z}\,x\,dz$

$= \int \frac{1}{z}\,dz$

© The Institute of Chartered Accountants of India

= $\log z$

= $\log (\log x) + c$

ans. (b)

167. Let $I = \int \log_{10}{}^{x}\, dx$

= $\int \log \frac{x}{e} \cdot \log_{10}^{e}\, dx$

= $\log_{10}^{e} \int \log x.1\, dx$

= $\log_{10}^{e} \left[\log x.x - \int \frac{1}{x} \times dx \right]$

$I = \log_{10}^{e} \left[x \log x - x \right] + c$

168. Let $I = \int \frac{4e^{x} + 6e^{-x}}{9e^{x} - 4e^{-x}}\, dx$

$\therefore \quad I = \int \frac{4e^{2x} + 6}{9e^{2x} - 4}\, dx$

Let $t = e^{2x}$

$\therefore dt = 2\ e^{2x}\, dx$

= $2t\ dx$

$\therefore I = \int \frac{4t + 6}{9t - 4} \frac{dt}{2t} = \int \frac{2t + 3}{t(9t - 4)} dt$

= $\int \left[\frac{0 + 3}{t(0 - 4)} + \frac{2(4/9) + 3}{4/9(9t - 4)} \right] dt$

= $\int \left[-\frac{3}{4t} + \frac{35}{4(9t - 4)} \right] dt$

= $-\frac{3}{4} \log t + \frac{35}{4} \cdot \frac{\log(9t - 4)}{9} + c$

= $-\frac{3}{4} \log e^{2x} + \frac{35}{36} \log \left(9e^{2x} - 4 \right) + c$

Ans. (a)

169. See formula from the text book

Ans. (c)

© The Institute of Chartered Accountants of India

170. Put $\sqrt{x^2-6x+100} = t$

$\therefore \quad x^2-6^x+100 = t^2$

$(2x-6)\,dx = 2t\,dt$

$(x-3)\,dx = t\,dt$

$$\int (x-3)\sqrt{x^2-6x+100}\,dx = \int \sqrt{x^2-6x+100}\,(x-3)dx$$

$$= \int t.tdt$$

$$= \int t^2\,dt$$

$$= \int \frac{t^3}{3}+c$$

$$= \frac{1}{3}(x^2-6^x+100)^{3/2}+c$$

Ans. (c)

171. No. of ways in which one or more friends may invited

$$= {}^6C_1 + {}^6C_2 + {}^6C_3 + {}^6C_4 + {}^6C_5 + {}^6C_6$$

$= 2^6 - 1 = 63$ ways.

Ans. (a) 63 ways.

172. No. of ways of failure of candidate.

$$= {}^4C_1 + {}^4C_2 + {}^4C_3 + {}^4C_4$$

$= 2^4 - 1 = 15$ ways.

Ans. (c) 15

© The Institute of Chartered Accountants of India

173. A voter can vote in the following ways

$254 = n_{C_1} + n_{C_2} + n_{C_3} + n_{C_4} + n_{C_5} + n_{C_6} + n_{C_7} .... n_{C_{n-1}}$

$\therefore 254 = 2^n - (n_{C_n} + 1) = 2^n - (1+1)$

$\therefore 256 = 2^n$

$\therefore 2^8 = 2^n \Rightarrow \therefore n = 8$ Total candidates = 8

Ans. (a) 8

174. No. of words of 3 consonants and 2 vowels among 17 consonants and 5 vowels are

$= 17_{C_3} \times 5_{C_2} \times 5!$

= 816000

Ans. (b) 81,6000

176. The present value of annual profit

V = A.P. (ni)

= 34000 × 3.7079

V = 128886 which is less than initial cost of machine. Machine must not be purchased

Ans. (a) Machine should not be purchased.

177. $40 = 2000 \times \frac{r}{100} \times 4$

$\therefore$ r = -0.5%

Ans. (b) 0.5%

© The Institute of Chartered Accountants of India

ANSWERS

178. $I_1 = 2000 \times \frac{4}{100} \times 1 = 80$

$I_2 = 3000 \times \frac{14}{100} \times 1 = 420$

Total Interest = 500

$\therefore$ rate of interest = $\frac{500 \times 100}{5000 \times 1} = 10\%$

r = 10%

Ans. (a) r = 10%

179. $I_1 - I_2 = 30 \Rightarrow 1200 \times \frac{R}{100} \times 3 - 1000 \frac{R}{100} \times 3 = 30$

$\Rightarrow 36R - 30R = 30$

$6R = 30$

$\therefore R = 5\%$

Ans. (c) 5%

180. $40 = 2000 \times \frac{2}{100} \times n$

n = 1 yr.

Ans. (a) = 1 yr.

181. $\frac{a+b}{2} = 20 \Rightarrow a + b = 40 \quad \rightarrow (1)$

SD = 5 $\rightarrow \frac{a-b}{2} = 5$

$\therefore$ a ... b = 10 $\rightarrow$ (2)

$\Rightarrow$ 2a = 50 $\Rightarrow$ a = 25

$\therefore$ b = 15

Ans. (a) 25, 15

© The Institute of Chartered Accountants of India

182. Mean $= \dfrac{\sum x}{n}$

$4.4 = \dfrac{1+2+6+a+b}{5}$

$\therefore \quad a + b = 13 \quad \rightarrow (1)$

$\sigma^2 = \dfrac{\sum x^2}{N} - \bar{x}^2$

$8.24 = \dfrac{\sum x^2}{5} - (4.4)^2$

$\sum x^2 = 138$

$1+4+9+a^2+b^2 = 138$

$\therefore \quad a^2 + b^2 = 138 \Rightarrow a^2 + (13 - a)^2 = 138$

$\Rightarrow \quad a^2 - 13a - 36 = 0 \qquad a = 9, 4$

$\therefore \quad$ Nos $\rightarrow 9, 4$

183. For individual series, the rank of the median is = $\left(\dfrac{N+1}{2}\right)^{th}$ term

Ans. (b) $\left(\dfrac{N+1}{2}\right)^{th}$ term

184. Rank of the median of the series 2, 3, 4, 5, 6, 7

$= \quad \left(\dfrac{N+1}{2}\right)^{th}$ term $= \left(\dfrac{6+1}{2}\right)^{th}$ term

$=$ 3.5 th term

Ans. (a) 3.5

185. Regression Eq. $2x + 3y - 10 = 0$

If $y = 50$

$\therefore 2x = +10 - 3 \times 50 = -140$

$x = -70$

None of these

Ans. (d) None of these.

© The Institute of Chartered Accountants of India

ANSWERS

186. Ans. (c) Refer Properties

187. Ans. (c) Refer Properties

188. Given r (x, y) = 0.4 $\rightarrow$ (1)

We know that r (a X, cY) = $\frac{a \times c}{|a| \times |c|}$.r(x,y) $\rightarrow$ (2)

Using (2) in (1), we get

r (2x, – y) = r (2x, – 1y)

= $\frac{2 \times (-1)}{|2| \times |-1|}$.r(x,y)

= $\frac{-2 \times 0.4}{2 \times 1}$

r (2x, – y) = – 0.4

Ans. (b)

189. Computation of Correlation Coefficient

	x	y	xy	x^2	y^2
	69	70	4830	4761	4900
	85	87	7395	7225	7569
Total	154	157	12225	11986	12469

$\bar{x} = \frac{154}{2} = 77, \quad \bar{y} = \frac{157}{2} = 78.5$

Cov (x,y) = $\frac{\sum x_i y_i}{n} - \bar{x}\,\bar{y} = \frac{12225}{2} - (77)\ (78.5)$

= 68

Sx = $\sqrt{\frac{\sum x_i^2}{n} - \bar{x}^2} = \sqrt{\frac{11986}{2} - (77)^2}$

= 8

Sy = $\sqrt{\frac{\sum y_i^2}{n} - \bar{y}^2} = \sqrt{\frac{12469}{2} - (78.5)^2} = 8.5$

$\therefore$ n = $\frac{\text{Cov}\ (x,y)}{\text{Sx}\ \text{Sy}} = \frac{68}{8 \times 8.5} = \frac{68}{68} = 1$

Ans. (a)

© The Institute of Chartered Accountants of India

190. Computation of correlation Co – efficient.

	x	y	xy	x^2	y^2
	102	50	5100	10404	2500
	109	48	5232	11881	2304
Total	211	98	10332	22285	4804

$$\bar{x} = \frac{211}{2} = 105.5,\ \bar{y} = \frac{98}{2} = 49$$

$$\text{Cov}(x, y) = \frac{\sum xi\, yi}{n} - \bar{x}\,\bar{y} = \frac{10332}{2} - (105.5)(49)$$

$$= \quad 5166 - 5169.5 = -3.5$$

$$Sx = \sqrt{\frac{\sum xi^2}{n} - \bar{x}^2} = \sqrt{\frac{22285}{2} - (105.5)^2} = 3.5$$

$$Sy = Sx = \sqrt{\frac{\sum yi^2}{n} - \bar{y}^2} = \sqrt{\frac{4804}{2} - (49)^2} = 1$$

$$\therefore \quad n = \frac{\text{Cov}(x, y)}{Sx\ Sy} = \frac{-3.5}{(3.5)\times(1)} = -1$$

Ans. (b)

191. Ans. (a) … Refer Properties

192. Ans. (a) … Refer Properties

193. Ans. (b) … Refer Properties

194. Given $X \sim N(\mu, \sigma 2)$, where $\mu = 2$ and $\sigma^2 = 9$

$\sigma = 3$

We want x so that

$P(2 \leq \times \leq x) = 0.4115 \rightarrow (1)$

When X = 2, $Z = \frac{x-\mu}{\sigma} = \frac{2-2}{3} = 0$

When x = x, $Z = \frac{x-2}{3}$ = Z1 (Say) $\rightarrow (2)$

From (1), we get $P(0 \leq Z \leq Z_1) = 0.4115$

$\Rightarrow \quad Z_1 = 1.35$ (from Normal Table)

© The Institute of Chartered Accountants of India

Substituting in (2), we get

$\frac{x-2}{3} = 1.35$

$\therefore$ x = 2+3 (1.35)

x = 6.05

Ans. (b)

195. Mean = First moment about origin = 35 (given) $\rightarrow$ (1)

Second moment about 35 = 10 (given)

$\Rightarrow$ Second moment about mean = 10

$\mu 2 = 0 \rightarrow (2)$

Since the given distribution is normal,

$\beta 1 = 0$ and $\beta 2 = 3$

$\therefore \beta 1 = \frac{\mu 3^2}{\mu 2^3} = 0 \Rightarrow \mu 3 = 0$

$\beta 2 = \frac{\mu 4}{\mu 2^2} = 3 \Rightarrow \mu 4 = 3\ \mu_2^2 = 3 \times 10^2 = 300$

$\therefore \mu 1 = 0$ (always), $\mu 2 = 10$, $\mu 3 = 0$, $\mu 4 = 300$

Ans. (c)

196. The most commonly used confidence limit is $\rightarrow$ 95%

Ans. (c) 95%

197. Sample mean is statistic

Ans. (b) Statistic

198. Deliberate sampling is – Non \random sampling

Ans. (b) Non random sampling

199. Stratified random sampling issued for Non – Homogeneous population.

Ans. (b) Non – homogeneous

200. Random Sampling is also called lottery sampling

Ans. True

© The Institute of Chartered Accountants of India

151. Let given number is x

Then the condition $\frac{1}{5}\left(\frac{1}{3}\left(\frac{1}{2}x\right)\right)=15$

$$\frac{x}{30}=15$$

$$x = 450$$

152. Let the number be = x.

Then given the condition = $\frac{3}{4}\left(\frac{1}{5}x\right)=60$

$$\frac{3x}{20}=60$$

$$x=\frac{60\times 20}{3}=400$$

Ans. (b)

153. Let the number be x.

Given $\frac{4}{5}\left[\frac{3}{8}(x)\right]=24$

$$\frac{3}{10}x=24$$

$$x=24\times\frac{10}{3}=80$$

$\therefore$ 250% of x = 250% of 80 = $\frac{250}{100}\times 80=200$

Ans. (d)

154. Let the number be x

Given condition $x+x^2=182$

$$x^2+x-182=0$$

$$(x+14)(x-13)=0$$

$$x=-14,\ x=13$$

$\therefore$ x = 13 (negative reflected)

Ans. (a)

© The Institute of Chartered Accountants of India

ANSWERS

155. Let the unit's digit of the number be x and ten's digit by y

Then $x + y = 12 \rightarrow (1)$

and the number = $10y + x$

Reversing the order of digits of the given number,

Unit's digit becomes y

and ten's digits becomes x

$\therefore$ New number = $10x + y$

According to the given condition of the problem $(10x + y) - (x + 10y) = 18$

$9x - 9y = -18$

$x - y = -2 \rightarrow (2)$

Adding (1) and (2) $\Rightarrow 2x = 10$

$x = 5$

$\therefore \quad \Rightarrow y = 7$

$\therefore$ The number is 75

Ans. (a)

156. Let the number of coins is x

Given $10x + \frac{14}{2}x + \frac{18}{4}x = 430$

$$\frac{40x + 28x + 18x}{4} = 430$$

$$\frac{86x}{4} = 430$$

$$x = \frac{430 \times 4}{86} = 20$$

$\therefore$ The one Rupee coins = $10x = 10 \times 20 = 200$

The 50 paise coins = $14x = 14 \times 20 = 280$

The 25 paise coins = $18x = 18 \times 20 = 360$

Ans. (a)

157. First Vessels Contain Milk Ratio 5

First Vessels Contain Water Ratio 2

© The Institute of Chartered Accountants of India

Second Vessels Contain Milk Ratio 6

Second Vessels Contain Water Ratio 1

Both the Vessels Milk = 5 + 6 = 11

Both the Vessels Water = 2 + 1 = 3

$\therefore$ The new Ratio = 11:3

Ans. (b)

158. Let the two numbers are x and y

Given $8x = 5y \rightarrow (1)$

and $x + 27 = y \rightarrow (2)$

$(2) \Rightarrow x = y - 27$

$(1) \Rightarrow 8(y - 27) = 5y$

$8y - 216 = 5y$

$3y = 216$

$y = \frac{216}{3} = 72$

$\therefore$ $x = 72 - 27 = 45$

$\therefore$ Sum of two number = x + y = 72 + 45 = 117

Ans. (c)

159. Let their monthly incomes be Rs. 9x and Rs. 7x respectively.

Let their monthly expenditures be Rs. 4y and Rs. 3y respectively.

According ot the given condition of the problem,

$9x - 4y = 200 \rightarrow (1)$

$7x - 3y = 200 \rightarrow (2)$

Multiply (1) by 3, we get

$27x - 12y = 600$

Multiply (2) by 4, we get

$28x - 12y = 800$

Subtracting (3) from (4), we get

$x = 200$

Hence their monthly income are Rs. (9×200=1800) and Rs. (7 × 200 = 1400).

Ans. (a)

© The Institute of Chartered Accountants of India

ANSWERS

160. Let x be the distributed amount of A, B and C

Given

$\therefore$ $5x + 11x + 3x = 950$

$19x = 950$

$x = \frac{950}{19} = 50$

$\therefore$ The amount of A = $5x = 5 \times 50 = 250$

The amount of B = $11x = 11 \times 50 = 550$

$\therefore$ The difference of A and B = 300

Ans. (a)

161. $\underset{x\to 1}{Lt} \frac{e^{-x} - e^{-1}}{x-1} \Rightarrow \underset{x\to 1}{Lt} \frac{e^{1-x} - 1}{e(x-1)}$

let $1 + h \to x$ where $h \to 0$

$\therefore$ $\underset{x\to 0}{Lt} \frac{e^{h} - 1}{e(-h)} = \frac{-1}{e}$

Ans. (b) $-1/e$

162. $\underset{x\to 0}{Lt} \frac{(1+x)^{n} - 1}{x}$

$= \underset{x\to 0}{Lt} \frac{(1 + nx + \frac{n(n-1)x^2}{2!} + ...) - 1}{x}$

$= \underset{x\to 0}{Lt} \frac{x\left[n + \frac{n(n-1)^{x}}{2!} + \frac{n(n-1)(n-2)x^2}{3!} .....\right]}{x}$

App Lt

n + 0 = n

Ans. (c) n

163. $\underset{x\to 0}{Lt} \frac{(x+2)^{5/3} - (a+2)^{5/3}}{x-a}$

$\underset{x\to 0}{Lt} \frac{(x+2)^{5/3} - (a+2)^{5/3}}{[(x+2)-(a+2)]}$

App Lt

© The Institute of Chartered Accountants of India

$\Rightarrow \frac{5}{3}.(a+2)^{5/3-1} = \frac{5}{3}(a+2)^{2/3}$

Ans. (a) $\frac{5}{3}(a+2)^{2/3}$

165. $\underset{x\to 0}{Lt}\ \frac{2^x-3^x}{x} \Rightarrow \underset{x\to 0}{Lt}\ \frac{(2^x-1)-(3^x-1)}{x}$

$\underset{x\to 0}{Lt}\ \frac{2^x-1}{x} - \underset{x\to 0}{Lt}\ \frac{3^x-1}{x} \qquad \left[\underset{x\to 0}{Lt}\ \frac{a^x-1}{x} = \log e^a\right]$

$\Rightarrow \log 2 - \log 3$

$\Rightarrow \log\left(\frac{2}{3}\right)$

Ans. (b)

166. $f'(x) = 3x^2 + 2$

$\int f'(x)\,dx = \int (3x^2+2)\,dx$

$f(x) + c = \frac{3x^3}{3} + 2x + c$

When $f(0) = 0 \Rightarrow c = 0$

$\therefore \quad f(x) = x^3 + 2x$

$\therefore \quad f(2) = 2^3 + 2(2)$

$= \quad 8 + 4 = 12$

Ans. (c)

167. Let $I = \int \frac{x+3}{x^2+6x+4}$

Put $x^2+6x+4 = t$

$\therefore (2x+6)dx = dt$

$(x+3)dx = \frac{dt}{2}$

$\therefore \int \frac{x+3}{x^2+6x+4}dx = \int \frac{dt}{2t}$

$= \frac{1}{2}\int \frac{1}{t}dt$

© The Institute of Chartered Accountants of India

$\frac{1}{2}\log(t)$

$\therefore \int \frac{x+3}{x^2+6x+4}dx = \frac{1}{2}\log(x^2+6x+4) + c$

168. $\int e^x \frac{x-1}{(x+1)^3}dx = \int \frac{x+1-2}{(x+1)^3}e^x dx$

$= \int e^x \left[\frac{1}{(x+1)^2} - \frac{2}{(x+1)^3}\right]dx$

$= \int e^x \{f(x)+f'(x)\}dx$

$= e^x f(x)$ where $f(x) = \frac{1}{(x+1)^2}$

$\int e^x \frac{(x-1)}{(x+1)^3}dx = \frac{e^x}{(x+1)^2} + c$

Ans.(a)

169. $\int (3x+5)^4 \, dx = \frac{(3x+5)^{4+1}}{(4+1)(3)} + c$

$= \frac{(3x+5)^5}{15} + c$

Ans. (b)

170. $\int \sqrt{7x+5} \, dx$

$= \int (7x+5)^{\frac{1}{2}} \, dx$

$= \frac{(7x+5)^{\frac{1}{2}+1}}{\left(\frac{1}{2}+1\right)(7)} + c$

$= \frac{(7x+5)^{\frac{3}{2}}}{\left(\frac{3}{2}\right)(7)} + c$

© The Institute of Chartered Accountants of India

$$\frac{2(7x+5)^{\frac{3}{2}}}{21}+c$$

171. Voter has option

(i) Two candidates from gentlemen = $3_{c_2}=3$

(ii) Two candidates from ladies = $3_{c_2}=3$

(iii) One from each ladies & gentlemen = $3_{c_1}\times 3_{C_1}=9$

Total options = 3 + 3 + 9 = 15

Ans. (c) 15

172. Total hand shakes in the party = $40_{c_2}=780$

Ans. (a) 780

173. Total triangle formed by m sides = m_{C_3}

$$\Rightarrow \quad \frac{m(m-1)(m-2)(m-3)!}{(m-3)!\,3!}$$

$$\Rightarrow \quad \frac{m(m-1)(m-2)}{6}$$

Ans. (a) $\frac{m(m-1)(m-2)}{6}$

174. Cricket team of 11 among 14 players out of which one wicketkeeper

$= \quad 12_{C_{10}} \times 2_{C_1} = 66\times 2$

$\Rightarrow$ 132

Ans. (b) 132

175. No. of ways in which a particular child goes to circus = $7_{C_2}\times 1 = 21$

Ans. (c) 21

176. $a^x = b^y = c^z = k$ (let)

$\therefore \log a^k = x,\ \log_b k = y,\ \log_c k = z$

$\therefore \log_k a = 1/x,\ \log_k b = 1/y,\ \log_k c = 1/z$

x, y, z in GP

$\therefore y^2 = xz$

$(\log_b k)^2 = (\log_a k).(\log_c k)$

© The Institute of Chartered Accountants of India

$\therefore \quad \frac{\log_b k}{\log_a k} = \frac{\log_c k}{\log_b k} \Rightarrow \frac{\log_k a}{\log_k k} = \frac{\log_k b}{\log_k c}$

$\therefore \log_a, \log_b$ and $\log_c$ in GP

Ans. (b) G.P.

177. $\frac{1}{1024} = 8.\left(\frac{1}{2}\right)^{n-1} \Rightarrow \left(\frac{1}{2}\right)^{13} = \left(\frac{1}{2}\right)^{n-1}$

$\therefore n = 14$

6th team from end = (14 – 6+1) from beginning = 9th term

$T_9 = 8.\left(\frac{1}{2}\right)^{9-1} = 8.\left(\frac{1}{2}\right)^{8} = \frac{1}{32}$

Ans. (c) 1/32

178. Product of 2nd term from start & last 2nd term from end = $(ar) \times a(r)^{n...2} = a^2 r^{(n...1)}$

Product of first & last term = $a \times ar^{n-1} = a^2 r^{n-1}$

Hence proved the statement. It is true statement

Ans. (a) True

179. a, b, c in GP $\therefore b^2 = ac$

a, x, b in AP $\Rightarrow x = \frac{a+b}{2}$

b, y, c in AP $\Rightarrow y = \frac{b+c}{2}$

$\therefore \frac{a}{x} + \frac{c}{y} = \frac{2a}{a+b} + \frac{2c}{b+c} \Rightarrow \frac{2[ab+ac+ac+bc]}{ab+b^2+ac+bc}$

$\Rightarrow \frac{2[ab+ac+ac+bc]}{ab+ac+ac+bc} \quad \{b^2 = ac\}$

$= 2$

Ans. (c) 2

180. $\frac{1}{x} + \frac{1}{y} = \frac{x+y}{xy} \Rightarrow \frac{\frac{a+b}{2} + \frac{b+c}{2}}{\frac{(a+b)(b+c)}{4}}$

$\Rightarrow \frac{2(a+2b+c)}{(ab+b^2+ac+bc)} = \frac{2(a+2b+c)}{ab+b^2+b^2+bc}$

© The Institute of Chartered Accountants of India

$= \frac{2(a+2b+c)}{b(a+2b+c)} = \frac{2}{b}$

Ans. (b) $\frac{2}{b}$

181. The number of times a particular item occurs in a given data is called its frequency.

Ans. (b) Frequency

182. Lower class (s) = 10.6

Width = 2.5 (Class interval)

Upper Class (L) of Lightest Class

$= S = 10 \times C.I.$

$= 10.6 + 10 \times 2.5$

$= 35.6$

Ans. (a) 35.6

183. $\frac{\text{Lower Class} + \text{Upper Class}}{2} = \text{Mid Value}$

$\frac{L + U.Class}{2} = m$

$\therefore$ Upper class = (2m – L)

Ans. (c) (2m – L)

184. Mean $= \frac{1.x + 2.2x + 3.3x + ...n.nx}{n(n+1).x/2}$

$= \frac{(1^2 + 2^2 + 3^2 + ...n^2)x}{n(n+1).x/2}$

$= \frac{(1^2 + 2^3 + 3^2 + ...n^2)x}{n(n+1).x/2}$

$= \frac{n(n+1)(2n+1)x \times 2}{6n(n+1) - x} = \frac{(2n+1)}{3}$

Mean $= \frac{(2n+1)}{3}$

Ans. (c) $\frac{2n+1}{3}$

© The Institute of Chartered Accountants of India

ANSWERS

185. $\bar{x} = 10, n = 4$

$\Sigma x = n.\ \bar{x} = 4 \times 10 = 40$

Corrected $\Sigma x = (40 + 4a)$

Corrected $\bar{x} = \frac{40 + 4a}{4}$

$13 = \frac{40 + 4a}{4}$

$4a = 12$

$a = 3$

Ans. (c) 3

186. From the given data, we observe that

$20 + 5 = 25$

$21 + 4 = 25$

and $22 + 3 = 25$

Thus, x and y are connected by the linear relation: $x + y = 25 \quad \rightarrow (1)$

$\Rightarrow$ There is perfect correlation between x and y

$\Rightarrow \quad r = \pm 1 \rightarrow (2)$

From (1) / We get $y = 25 - x$

$\therefore$ As x increases, y decreases (by the same amount)

$\Rightarrow$ x and y are negatively correlated $\rightarrow (3)$

From (2) and (3), we conclude that

$r = r(x, y) = -1$

Ans. (c)

187. Ans. (b) Refer Properties

188. Ans. (b) Refer Properties

189. Ans. (b) Refer Properties

190. Ans. (b) Refer Properties

191. Lt X … B (n=6, p). When X denotes the number of successes. Then, by binomial probability law, the probability of r successes is givenby

$p(r) = P(x=r) = 6C_r P^r q^{6-r} \rightarrow (1)$

r = 0, 1, 2, ... 6

Put r = 3 and 4 in (1)

© The Institute of Chartered Accountants of India

(1) $\Rightarrow p(3) = 6C_3 p^3 q^3 = 20p^3 q^3 = 0.2457$ (given)

$p(4) = 6C_4 p^4 q^2 = 15 p^4 q^2 = 0.0819$ (given)

$$\frac{p(4)}{p(3)} = \frac{15p^4q^2}{20p^3q^3} = \frac{0.0819}{0.2457} = \frac{1}{3}$$

$$\Rightarrow \quad \frac{3}{4}.\frac{p}{q} = \frac{1}{3}$$

$\therefore \quad 9p = 4q = 4(1 - p)$

$\therefore \quad 13p = 4$

$p = 4/13$

$\therefore \quad q = 1 - p = 1 - 4/13 = 9/13$

192. Ans. (a) – Refer Properties

193. Ans. (b) – Refer Properties

194. Ans. (a) – Refer Properties

195. Ans. (b) – Refer Properties

196. Which measure of dispersion has some desirable mathematical properties → Standard Deviation.

Ans. (a) Standard Deviation.

197. $x\sqrt{1+y} + y\sqrt{1+x} = 0$

$$\Rightarrow x\sqrt{1+y} = -y\sqrt{1+x}$$

Eg. Both side

$x^2(1+y) = y^2(1+x)$

$(x^2 - y^2) = y^2x - x^2y$

$(x+y)(x-y) = -xy(x-y)$

$\therefore \quad x + y + xy = 0 \quad \Rightarrow y = \frac{-x}{1+x}$

$$\therefore \quad \frac{dy}{dx} = \frac{(1+x)(-1) - (-x)(1)}{(1+x)^2} = \frac{-1-x+x}{(1+x)^2}$$

$$\therefore \quad \frac{dy}{dx} = \frac{-1}{(1+x)^2} \Rightarrow (1+x^2)\frac{dy}{dx} = -1$$

Ans. (c) (–1)

© The Institute of Chartered Accountants of India

ANSWERS

198. $y = x\sqrt{x^2+1} + \log\left(x+\sqrt{x^2+1}\right)$

$$\frac{dy}{dx} = x.\frac{1}{2\sqrt{x^2+1}}.2x + \sqrt{x^2+1}.1 + \frac{1}{x+\sqrt{x^2+1}}\left(1+\frac{1}{2\sqrt{x^2+1}}.2x\right)$$

$$= \frac{x^2}{\sqrt{x^2+1}} + \sqrt{x^2+1} + \frac{\sqrt{x^2+1}+x}{\left(x+\sqrt{x^2+1}\right)}\frac{1}{\sqrt{x^2+1}}$$

$$= \frac{x^2+x^2+1+1}{\sqrt{x^2+1}} = \frac{2(x^2+1)}{\sqrt{x^2+1}} = 2\sqrt{1+x^2}$$

Ans. (c) $2\sqrt{1+x^2}$

199. $y = ae^{mx} + be^{-mx}$

$$\therefore \quad \frac{dy}{dx} = ame^{mx} - bme^{-mx}$$

$$\frac{d^2y}{dx^2} = \frac{d}{dx} = \left(ame^{mx} - bme^{-mx}\right)$$

$$= am^2e^{mx} + bm^2e^{-mx}$$

$$= m^2\left(ae^{mx} + be^{-mx}\right)$$

$$\frac{d^2y}{dx^2} = m^2y$$

Ans. (c) m^2y

200. $12_{C_5} + 2.12_{C_4} + 12_{C_3} = 14_{C_x}$

$$\left(12_{C_5} + 12_{C_4}\right) + \left(12_{C_4} + 12_{C_3}\right) = 14_{C_x}$$

$$13_{C_5} + 13_{C_4} = 14_{C_x}$$

$$\left[{}^nC_r + {}^nC_{r-1} = {}^{n+1}C_r\right]$$

$$14_{C_5} = 14_{C_x}$$

x = 5 but value $14_{C_9} = 14_{C_5}$

$\therefore$ $14_{C_x} = 14_{C_9}$

$\therefore$ x = 9

© The Institute of Chartered Accountants of India

x = 5 or 9

Ans. (c) 5 or 9.

Model Test Paper – BOS/CPT – 7

151. Let the number be x. Then according to the given condition of the problem,

$$\frac{x}{3}=\frac{x+1}{4}+1$$

$$\Rightarrow \frac{x}{3}-\frac{x+1}{4}=1$$

$$\Rightarrow \frac{4x-3(x+1)}{12}=1$$

$$\Rightarrow \frac{x-3}{12}=1$$

$\Rightarrow x - 3 = 12$

$\Rightarrow x = 3 + 12 = 15$

Hence the required number is 15

152. Let the fraction = x

And Correct answer = y

$\therefore$ Given $\frac{16}{17}x = y \rightarrow (1)$

and $\frac{x}{\frac{16}{17}} = y + \frac{33}{340}$

i.e. $\frac{17}{16}x = y + \frac{33}{340} \rightarrow (2)$

(1) $\Rightarrow x = \frac{17}{16}y$

Substitute x the value of x in equation (2)

$$\frac{17}{16}\times\frac{17}{16}y = y+\frac{33}{340}$$

$$\frac{289}{256}y = y + \frac{33}{340}$$

© The Institute of Chartered Accountants of India

$$\frac{289}{256}y - y = \frac{33}{340}$$

$$\left(\frac{289-256}{256}\right)y = \frac{33}{340}$$

$$\frac{33}{256}y = \frac{33}{340}$$

$$y = \frac{33}{340} \times \frac{256}{33}$$

$$y = \frac{64}{85}$$

Ans. (a)

153. Let the number be x

Given $\frac{5}{7}\left[\frac{4}{15}(x)\right] = 8 + \frac{2}{5}\left[\frac{4}{9}(x)\right]$

$$\frac{4}{21}x = 8 + \frac{8}{45}x$$

$$\frac{4}{21}x - \frac{8}{45}x = 8$$

$$\left(\frac{180-168}{945}\right)x = 8$$

$$\frac{12}{945}x = 8$$

$$x = 8 \times \frac{945}{12}$$

$$x = 708.75$$

Ans. (d)

154. Let two numbers are x and y

Given condition $x + y = 14 \rightarrow (1)$

$y - x = 10 \rightarrow (2)$

Adding (1) and (2) $\Rightarrow 2x = 24$

$x = 12$

(1) $\Rightarrow 12 + y = 14$

© The Institute of Chartered Accountants of India

$y = 2$

$\therefore$ Product of two numbers = $x \times y = 12 \times 2 = 24$

Ans. (a)

155. Let two numbers are x and y.

Given $x - y = 11 \rightarrow (1)$

And $\frac{x+y}{5} = 9$

i.e. $x + y = 45 \rightarrow (2)$

Adding (1) and (2) $\Rightarrow 2x = 56$

$X = 28$

(1) $\Rightarrow 28 - y = 11$

$-y = 11 - 28$

$= -17$

$y = 17$

$\therefore$ The two numbers are 28, 17.

Ans. (d)

156. Sub duplicate Ratio of 16:49 = $\sqrt{16} : \sqrt{49}$

= 4 : 7

Ans. (a)

157. Duplicate Ratio of 4 : 5 = $4^2 : 5^2$

= 16 : 25

Ans. (a)

158. Triplicate Ratio of 3 : 5 = $3^3 : 5^3$

= 27 : 125

Ans. (a)

159. The sub – triplicate Ratio of 8 : 125 = $\sqrt[3]{8} : \sqrt[3]{125}$

= 2 : 5

Ans. (b)

160. 4 th Proportion of 6, 8 and 15 is $\frac{6}{8} = \frac{15}{x}$

$6x = 15 \times 8$

© The Institute of Chartered Accountants of India

$x = \frac{15 \times 8}{6}$

$= 20$

Ans. (c)

161. Let the two numbers be x and y. According to the First condition of the problem,

$\frac{x}{y} = \frac{4}{1}$

$\Rightarrow \quad x = 4y \qquad \rightarrow (1)$

According to the second condition of the problem,

$\frac{x+5}{y+5} = \frac{3}{1}$

$x + 5 = 3 (y + 5)$

$x + 5 = 3y + 15$

$x - 3y = 15 - 5 = 10 \rightarrow (2)$

Put $x = 4y$ from (1) in (2), we get

$4y - 3y = 10$

$y = 10$

$(1) \Rightarrow x = 4 (10) = 40$

Hence the required numbers are 40 and 10.

Ans. (b)

162. Let A having money = 3x

B having money = 4x

C having money = 5x

Given $3x = 300$

$x = 100$

$\therefore \quad C = 5x = 5 \times 100 = 500$

Ans. (c)

163. Let the two numbers be x and y. According to the first condition of the problem.

$\frac{x}{y} = \frac{5}{6}$

$6x = 5y$

$6x - 5y = 0 \qquad \rightarrow (1)$

© The Institute of Chartered Accountants of India



According to the second condition of the problem

$$\frac{x-5}{y-5}=\frac{4}{5}$$

$5(x-5) = 4(y-5)$

$5x - 25 = 4y - 20$

$5x - 4y = 5 \quad \rightarrow (2)$

$(1) \times 5 \Rightarrow 30x - 25y = 0 \quad \rightarrow (3)$

$(2) \times 5 \Rightarrow 30x - 24y = 30$

Subtracting (3) from (4), we get

$y = 30$

$(1) \Rightarrow 6x - 5(30) = 0$

$6x = 150$

$x = \frac{150}{6} = 25$

Hence the required numbers are 25 and 30

Ans. (c)

164. Let the given numbers be x and y. Then according to the given conditions of the problem.

$$\frac{x+1}{y+1}=\frac{1}{2}$$

$\Rightarrow \quad 2x + 2 = y + 1$

$2x - y = -1 \quad \rightarrow (1)$

and $\frac{x-5}{y-5}=\frac{5}{11}$

$11x - 55 = 5y - 25$

$11x - 5y = 30 \quad \rightarrow (2)$

$\therefore \quad (1) \times 11 \Rightarrow 22x - 11y = -11 \quad \rightarrow (3)$

$(2) \times 2 \Rightarrow 22x - 10y = 60 \quad \rightarrow (4)$

$(3) - (4) \Rightarrow -y = -71$

$y = 71$

$(1) \Rightarrow 2x - 71 = -1$

$2x = 70$

© The Institute of Chartered Accountants of India

$x = 35$

Hence, the required numbers are 35 and 71

Ans. (b)

165. Let the number to be subtracted be x.

Then according to the problem

$$\frac{27-x}{43-x}=\frac{7}{15}$$

$\Rightarrow$ $15(27-x) = 7(43-x)$

$\Rightarrow$ $405 - 15x = 301 - 7x$

$15x - 7x = 405 - 301$

$8x = 104$

$$x = \frac{104}{8} = 13$$

Hence the required number is 13

Ans. (a)

166. Let the unit digit = x

and ten digits = y

$\therefore$ $x+y = 3 \quad \rightarrow (1)$

and the number = $10y + x$

Reversing the order of digits

Units digit = y

and ten's digit = x

$\therefore$ Number = $10x + y$

According to the given condition of the problem

$7(10y + x) = 4(10x+y)$

$70y + 7x = 40x + 4y$

$70x - 40x + 70y - 4y = 0$

$-33x + 66y = 0$

$-x+2y = 0 \quad \rightarrow 92)$

Adding (1) and (2) we get

$3y = 3$

$y = 1$

© The Institute of Chartered Accountants of India

(1) $\Rightarrow$ x + 1 = 3

x = 3 –1 = 2

Hence, the required number is 12

167. The committee of six must include atleast 2 ladies.

i.e. two or more ladies. As there are only 3 ladies, the following possibilities arise:

The committee of 6 consists of

(i) 4 men and 2 ladies, (ii) 3 men and 3 ladies.

The number of ways for (i) = $^{7}C_4 + {}^{3}C_2$

= $35 \times 3 = 105$

The number of ways for (ii) = $^{7}C_3 \times {}^{3}C_3$

= 35×1

= 35

Hence the total number of ways of forming a committee so as to include atleast two ladies = 105 + 35 = 140

Ans. (a)

168. We have $nCr = \frac{n!}{r!(n-4)!}$

Now substituting for n and r, we get

$^{28}C_{2r} = \frac{28!}{(2r)!(28-2r)!}$

$24C_{2r-4} = \frac{24!}{(2r-4)!\{24-(2r-4)\}!}$

$= \frac{24!}{(2r-4)!\,(28-2r)!}$

Given $28_{C_{2r}} : 24_{C_{2r-4}} = 225:11$

$\Rightarrow \frac{28_{C_{2r}}}{24_{C_{2r-4}}} = \frac{28!}{(2r)!(28-2r)!} \div \frac{(2r-4)!\,(28-2r)!}{24!}$

$= \frac{28\times27\times26\times25\times24!}{2r(2r-1)(2r-2)(2r-3)(2r-4)!\,(28-2r)!} \times \frac{(2r-4)!\,(28-2r)!}{24!}$

$= \frac{28\times27\times26\times25}{2r(2r-1)(2r-2)(2r-3)} = \frac{225}{11}$

© The Institute of Chartered Accountants of India

$\Rightarrow$ $2r(2r-1)(2r-2)(2r-3) = \frac{11\times28\times27\times26\times25}{225}$

$= 11 \times 28 \times 3 \times 26$

$= 11 \times 7 \times 4 \times 3 \times 13 \times 2$

$= 11 \times 12 \times 13 \times 14$

$= 14 \times 13 \times 12 \times 11$

$\therefore$ $2r = 14$

$r = 7$

Ans. (b)

169. Let in the number unit's digit = x

and ten's digit = y

$\therefore$ Number = $10y + x$

According to the given conditions of the problem,

$8(x+y) + 1 = 10y + x$

(or) $8x + 8y + 1 = 10y + x$

$\Rightarrow$ $8x - x + 8y - 10y + 1 = 0$

$7x - 2y + 1 = 0 \rightarrow (1)$

and $13(y - x) + 2 = 10y + x$

$13y - 13x + 2 = 10y + x$

$\Rightarrow$ $x + 13x + 10y - 13y - 2 = 0$

$\Rightarrow$ $14x - 3y - 2 = 0 \rightarrow (2)$

$(1) \times 2 \Rightarrow 14x - 4y + 2 = 0 \rightarrow (3)$

$(2) - (3) \Rightarrow y - 4 = 0$

$y = 4$

Put $y = 4$ in (1), we get

$7x - 8 + 1 = 0$

$7x = 7$

$x = 1$

Hence, the required number is 41

Ans. (b)

© The Institute of Chartered Accountants of India

170. We want to find out the number of combination of 12 things taken 3 at a time and this is by:

$$12_{C_3} = \frac{12!}{3!(12-3)!}$$

$$= \frac{12!}{3!\,9!} = \frac{12\times11\times10\times9!}{3!\,9!}$$

$$= \frac{12\times11\times10}{3\times2}$$

$$= 220$$

Ans. (c)

172. $$\lim_{x\to9} \frac{\sqrt{x}-3}{x-9} = \lim_{x\to9} \frac{\sqrt{x}-3}{\sqrt{x}-3}\frac{1}{\sqrt{x}+3}$$

$$\text{App lt} = \frac{1}{3+3} = \frac{1}{6}$$

Ans. (a) $\frac{1}{6}$

173. $$\lim_{x\to9} \frac{\sqrt{x+0}-\sqrt{2a}}{x-0} \Rightarrow \lim_{x\to9} \frac{\sqrt{x+a}-\sqrt{2a}}{x-0} \times \frac{\left(\sqrt{x+a}-\sqrt{2a}\right)}{\left(\sqrt{x+a}+\sqrt{2a}\right)}$$

$$\Rightarrow \lim_{x\to9} \frac{x+a-2a}{(x-a)\left(\sqrt{x+a}+\sqrt{2a}\right)} = \lim_{x\to a} \frac{(x-a)}{(x-a)\left(\sqrt{x+a}+\sqrt{2a}\right)}$$

App Lt

$$\frac{1}{\left(\sqrt{a+a}+\sqrt{2a}\right)} = \frac{1}{2\sqrt{2a}}$$

Ans. (b) $\frac{1}{2\sqrt{2a}}$

174. $$\lim_{x\to\infty} \frac{6+5x^2}{4x+15x^2} \Rightarrow \lim_{x\to\infty} x^2 \frac{\left(5+\frac{6}{x^2}\right)}{x^2\left(15+\frac{4}{x}\right)}$$

App Lt.

$$\Rightarrow \frac{5+0}{15+0} = \frac{1}{3}$$

Ans. (c) $\frac{1}{3}$

© The Institute of Chartered Accountants of India

ANSWERS

175. $\underset{x\to\infty}{Lt}\ \frac{a-bx}{x^2} \Rightarrow \underset{x\to\infty}{Lt}\left(\frac{a}{x^2}-\frac{b}{x}\right)$

App Lt.

$\Rightarrow (0 \ldots 0) = 0$

Ans.(a) 0

176. $y=\frac{e^x-e^{-x}}{e^x+e^{-x}} \Rightarrow \frac{dy}{dx} = \frac{(e^x+e^{-x})\frac{d}{dx}(e^x-e^{-x})-(e^x-e^{-x}).\frac{d}{dx}(e^x+e^{-x})}{(e^x+e^{-x})^2}$

$= \frac{(e^x+e^{-x})(e^x+e^{-x})-(e^x-e^{-x})(e^x-e^{-x})}{(e^x+e^{-x})^2}$

$= \frac{e^{2x}+e^{-2x}+2-e^{2x}-e^{-2x}+2}{(e^x+e^{-x})^2} = \frac{4}{(e^x+e^{-x})^2}$

Ans. (b) $\frac{4}{(e^x+e^{-x})^2}$

177. $y = \frac{x}{(1+x)^2} \Rightarrow \frac{dy}{dx} = \frac{(1+x)^2\frac{d}{dx}x-x\frac{d}{dx}(1+x)^2}{(1+x)^4}$

$\Rightarrow \frac{(1+x)^2.1-x.2(1+x)}{(1+x)^4} = \frac{1+x^2+2x-2x-2x^2}{(1+x)^4}$

$= \frac{1-x^2}{(1+x)^4} = \frac{1-x}{(1+x)^3}$

Ans. (b) $\frac{1-x}{(1+x)^3}$

178. $y = \sqrt{x+\sqrt{x}}$

$\frac{dy}{dx}=\frac{d}{dt}t^{1/2}.\frac{d}{dx}(x+\sqrt{x})$

$= \frac{1}{2\left(\sqrt{x+\sqrt{x}}\right)}.\left(1+\frac{1}{2\sqrt{x}}\right)$

© The Institute of Chartered Accountants of India

$$= \frac{2\sqrt{x}+1}{4\sqrt{x}\sqrt{x+\sqrt{x}}}$$

Ans. (a) $\frac{2\sqrt{x}+1}{4\sqrt{x}\sqrt{x+\sqrt{x}}}$

179. $y = 7^{x^2+2x}$

$$\frac{dy}{dx} = \frac{d}{dt}7^t.\frac{d}{dx}(x^2+2x)$$

$$= 7^{x^2+2x}.\log 7.(2x+2)$$

$$\frac{dy}{dx} = 2(x+1).7^{x^2+2}.\log 7$$

Ans. (b) $2(x+1).7^{x^2+2}.\log 7$

180. $y = \log\left(x+\sqrt{x^2+a^2}\right)$

$$\frac{dy}{dx} = \frac{d}{dt}\log t.\frac{d}{dx}\left(x+\sqrt{x^2+a^2}\right)$$

$$= \frac{1}{x+\sqrt{x^2+a^2}}.\left[1+\frac{1}{2\sqrt{x^2+a^2}}.2x\right]$$

$$= \frac{\left(\sqrt{x^2+a^2}+x\right)}{\sqrt{x^2+a^2}\left(x+\sqrt{x^2+a^2}\right)}$$

$$\frac{dy}{dx} = \frac{1}{\sqrt{x^2+a^2}}$$

Ans. (a) $\frac{1}{\sqrt{x^2+a^2}}$

181. Given $(x-y)e^{\frac{x}{x-y}} = a$

Differentiate on both sides.

© The Institute of Chartered Accountants of India

$$(x-y).e^{\frac{x}{x-y}}\left[\frac{(x-y)(1)-x.\left(1-\frac{dy}{dx}\right)}{(x-y)^2}\right]+e^{\frac{x}{x-y}}\left(1-\frac{dy}{dx}\right)=0$$

$$e^{\frac{x}{x-y}}\left[\frac{x-y-x+x\frac{dy}{dx}}{(x-y)}+1-\frac{dy}{dx}\right]=0$$

$$\frac{-y+x\frac{dy}{dx}}{(x-y)}+1-\frac{dy}{dx}=0$$

$$\frac{-y}{x-y}+\frac{x}{x-y}\frac{dy}{dx}+1-\frac{dy}{dx}=0$$

$$\frac{dy}{dx}\left[\frac{x}{x-y}-1\right]=\frac{y}{x-y}-1$$

$$\frac{dy}{dx}\left[\frac{x-x+y}{x-y}\right]=\frac{y-x+y}{x-y}$$

$$\frac{dy}{dx}\left[\frac{y}{x-y}\right]=\frac{2y-x}{x-y}$$

$$\frac{dy}{dx}=\frac{2y-x}{y}$$

$$\therefore\quad y\frac{dy}{dx}+x=y\left[\frac{2y-x}{y}\right]+x$$

$= \quad 2y - x + x$

$$y\frac{dy}{dx}+x=2y$$

Ans. (c)

182. Given demand law $x = \sqrt{10-p^2}$

$$x = (10-p^2)^{1/2}$$

$$\frac{dx}{dp}=\frac{1}{2}(10-p^2)^{\frac{1}{2}-1}(-2p)$$

© The Institute of Chartered Accountants of India

$$\frac{1(-2p)}{2\sqrt{10-p^2}}$$

$$\frac{dx}{dp} = \frac{-p}{\sqrt{10-p^2}}$$

$$|ed| = \frac{p}{x} \cdot \frac{dx}{dp}$$

$$= \frac{p}{(10-p)^{1/2}} \cdot \frac{-p}{2\sqrt{10-p^2}}$$

when p = 2

$$|ed| = \frac{2}{\sqrt{6}} \cdot \frac{-2}{\sqrt{6}}$$

$ed = -2/3$

$$|ed| = \frac{2}{3}$$

Ans. (a)

Alternate

183. $\therefore \int \frac{x^3}{x+1} dx$

$$= \int \left(x^3 - x + 1 - \frac{1}{x+1} \right) dx$$

$$= \frac{x^3}{3} - \frac{x^2}{2} + x - \log(x+1) + c$$

Ans. (c)

184. $\int \left(\frac{e^{4x} + e^{2x}}{e^{3x}} \right) dx$

$$= \int \left(\frac{e^{4x}}{e^{3x}} + \frac{e^{2x}}{e^{3x}} \right) dx$$

$$= \int e^x dx + \int e^{-x} dx$$

$= e^x - e^{-x} + c$

Ans. (b)

© The Institute of Chartered Accountants of India

ANSWERS

185. $\int \frac{x^4+1}{x^2+1}\,dx = \int\left(x^2-1+\frac{2}{x^2+1}\right)dx$

$= \int x^2\,dx - \int dx + 2\int \frac{1}{x^2+1}\,dx$

$= \frac{x^3}{3} - x + 2\tan^{-1} x + c$

Ans. (c)

186. Let $I = \int \log(x+1)\,dx$

$\therefore \quad I = \int \log(x+1).1\,dx$

Integrating by parts

[Here log (x+1) is to be taken as first function and unity as second function]

I = [log (x+1] integral of '1' – integral of

[d/dx (log (x+1)] + integral of '1']

$= \log(x+1).x - \int \frac{1}{x+1}.x\,dx$

$= x\log(x+1) - \int \frac{x+1-1}{x+1}dx$

$= x\log(x+1) - \int\left(1-\frac{1}{x+1}\right)dx$

$= x\log(x+1) - [x - \log(x+1)]$

$I = x\log(x+1) - x + \log(x+1) + c$

187. Consider $\frac{1}{\sqrt{x}+\sqrt{1+x}} = \frac{\sqrt{x}-\sqrt{1+x}}{x-(1+x)}$

$= \sqrt{1+x} - \sqrt{x}$

$\therefore \quad I = \int \frac{dx}{\sqrt{x}+\sqrt{1+x}}$

$= \int \sqrt{1+x}\,dx - \int \sqrt{x}\,dx$

$I = I_1 - I_2$

$I_1 = \int \sqrt{1+x}\,dx$

© The Institute of Chartered Accountants of India

Let $z = 1 + x$

$dz = dx$

$\therefore I_1 = \int\sqrt{1+x}\ dx = \int\sqrt{z}\ dz$

$= \frac{2}{3}z^{3/2}$

$= \frac{2}{3}(1+x)^{3/2}$

$\therefore I = 2/3\ (1+x)^{3/2} - \frac{x^{3/2}}{3/2} + c$

$I = 2/3\ \left\{(1+x)^{3/2} - x^{3/2}\right\} + c$

188. Consider $x^3 + x^2 - 2x = x(x^2 + x - 2)$

$= x(x^2 + 2x - x - 2)$

$= x\ \{x(x+2) - (x+2)\}$

$= x\ (x - 1)\ (x + 2)$

$\therefore$ We may write $\frac{x^2 - x + 2}{x^3 + x^2 - 2x} = \frac{x^2 - x + 2}{x(x-1)\ (x+2)}$

Let $\frac{x^2 - x + 2}{x(x-1)(x+2)} = \frac{A}{x} + \frac{B}{x-1} + \frac{C}{x+2}$

(or) $x^2 - x + 2 = A(x-1)\ (x+2) + Bx(x+2) + Cx(x-1)$

Substituting x = 1, We find 2 = 3B

B = 2/3

Substituting x = – 2, We find 8 = 6c

i.e. C = 4/3

Substituting x = 0, We find 2 = – 2A

A = – 1

$\therefore I = \int\frac{x^2 - x + 2}{x^3 + x^2 - 2x}dx = -\int\frac{dx}{x} + \frac{2}{3}\int\frac{dx}{x-1} + \frac{4}{3}\int\frac{dx}{x+2}$

I = – log x + 2/3 log (x – 1) + 4/3 log (x+2) + log c

Ans. (c)

© The Institute of Chartered Accountants of India

189. Let $I = \int \frac{1}{3x^2+13x-10}dx$

$= \frac{1}{3}\int \frac{1}{x^2+\frac{13x}{3}-\frac{10}{3}}dx$

$= \frac{1}{3}\int \frac{1}{\left[x^2+2.\frac{13}{6}x+\left(\frac{13}{6}\right)^2\right]-\left(\frac{13}{6}\right)^2-\frac{10}{3}}dx$

$= \frac{1}{3}\int \frac{1}{(x+13/6)^2-\frac{289}{36}}dx$

Let $t = x+13/6$

$\therefore dt = dx$

$\therefore I = 1/3 \int \frac{1}{t^2-(17/6)^2}dt$

$= \frac{1}{3}.\frac{1}{2(17/6)}\log\left[\frac{t-17/6}{t+17/6}\right]$

$= \frac{1}{17}\log\left[\frac{6t-17}{6t+17}\right]$

$= \frac{1}{17}\log\left[\frac{6(x+13/6)-17}{6\left(x+\frac{13}{6}\right)+17}\right]$

$= \frac{1}{17}\log\left[\frac{3x-2}{3x+15}\right]+c$

Ans. (b)

190. $\int e^x\{f(x)+f'(x)\}dx$

By the method of integration by parts, we may write

$\int e^x f(x)dx = f(x)\int e^x dx - \int\left\{\frac{d}{dx}f(x)\int ex\,dx\right\}dx$

$= e^x f(x) - \int e^x f'(x)\,dx$

© The Institute of Chartered Accountants of India

Transposing $\int e^x f(x)\,dx + \int e^x f'(x)\,dx = e^x f(x)$

(or) $\int e^x \{f(x) + f'(x)\}\,dx = e^x f(x)$

191. $\int_a^b \frac{\log x}{x} dx$ Let $\log x = t$ $x = a, t = \log a$

$$\frac{1}{x} = \frac{dt}{dx} \qquad x = b, t = \log b$$

$\int_{\log b}^{\log a} t.dt. \Rightarrow \left[\frac{t^2}{2}\right]_{\log a}^{\log b}$

$\Rightarrow \frac{1}{2}\left[(\log b)^2 - (\log a)^2\right] \Rightarrow \frac{1}{2}\left[\log(ab).\log\left(\frac{b}{a}\right)\right]$

Ans. (a) $\frac{1}{2}\left[\log(ab).\log\left(\frac{b}{a}\right)\right]$

192. $\int [f(x) + f(-x)][g(x) - g(-x)]\,dx$

$\Rightarrow \int 0.[g(x) - g(-x)]dx \Rightarrow 0$

Ans. (a) 0

193. $\int_a^b \frac{dx}{(a+b-x)^{2/3}}$ Let $a + b - x = t$ $x = a, t = b$

$$-1 = \frac{dt}{dx} \qquad x = b, t = a$$

$\Rightarrow -\int_b^a t^{-2/3}\,dt \Rightarrow \int_a^b t^{-2/3}\,dt$

$\Rightarrow 3\left[t^{1/3}\right]_a^b \Rightarrow 3\left[b^{1/3} - a^{1/3}\right]$

Ans. (a) $3.\left[b^{1/3} - a^{1/3}\right]$

194. $I = \int_0^2 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{2-x}} dx$ ... (i)

$I = \int_0^2 \frac{\sqrt{2-x}}{\sqrt{2-x} + \sqrt{x}}$ ... (ii) $[f(x) = f(a-x)]$

© The Institute of Chartered Accountants of India

ANSWERS

$2I = \int_0^2 dx = [x]_0^2 = 2$

$\therefore \quad I = \frac{1}{2} \times 2 = 1$

Ans. (a) 1

195. $I = \int_0^1 \log\left(\frac{1}{x} - 1\right)dx \Rightarrow \int_0^1 \log\left(\frac{1-x}{x}\right)dx$ ... (i)

$I = \int_0^1 \log\left(\frac{1-1+x}{1-x}\right)dx \qquad [f(x) = f(a-x)]$

$I = \int_0^1 \log \frac{x}{1-x} \Rightarrow -\int_0^1 \log\left(\frac{1-x}{x}\right) dx$ ... (ii)

$2I = 0 \qquad \therefore I = 0$

Ans. (c) 0

196. No. of ways in which 7 dept distributed among 3 minister

$= \left(7_{C_3} \times 4_{C_3} \times 1\right) + \left(7_{C_3} \times 4_{C2}\, x\right) \times 3!$

$= (120 + 330) \times 6 = 1980$

Ans. (d) None of these

197. No. of selections of letters

(i) 2 like and 1 different = $3_{C_1} \times 2_{C_1} = 3 \times 2 = 6$

(ii) 3 different = $5_{C_3} = 10$

$\therefore$ Total no. of ways of selection letters = 16

$\therefore$ Total words = 16×3! = 96 words.

Ans. (b) 96

198. No. of ways to form three digit nos. by using (1, 2, 3, 4, 3, 2) are – 42.

Ans. (b) 42

199. $S_n + S_{n-2} - 2.S_{n-1} = \frac{n}{2}[2a + (n-1)d] + \frac{n-2}{2}[2a+(n-2-1)d]$

$\ldots 2.\frac{n-1}{2}[2a + (n-1-1)d]$

© The Institute of Chartered Accountants of India

$= \frac{n}{2}[2a+(n-1)d] + \left(\frac{n-2}{2}\right)[2a+(n-3)d] - (n-1)[2a+(n-2)d]$

$= an + \frac{(n-1)n}{2}d + a(n-2) + \frac{(n-3)(n-2)}{2}d - 2a(n-1) - (n-1)(n-2)d$

$= d\left[\frac{(n-1)}{2}n + \frac{(n-3)(n-2)}{2} - (n-1)(n-2)\right]$

$d\left(\frac{n^2-n+n^2-5n^2+6-2n^2-4+6n}{2}\right)$

$= \frac{2d}{2} = d$

Ans. (a) d

200. $\frac{A3}{A(n-1)} = \frac{1}{3}$

$\frac{a+3d}{a+(n-1)d} = \frac{1}{3} \Rightarrow (3+3\ d)\ 3 = 3 + (n-1)d$

$\therefore \quad d = \frac{6}{n-10}$

$\therefore \quad 7n = a + (N...1)\ d \Rightarrow 31 = 3 + (n+2-1)\ \frac{6}{n-10}$

$28 = (n+1)\ \frac{6}{n-10} \quad \therefore n = 13$

Ans. (c) 13

Model Test Paper – BOS/CPT – 8

151. The first no. divide by 8 between 100 and 200 is 104

The last no. divide by 8 between 100 and 200 is 200

$\therefore$ The total number divide by 104 and 200 is 13.

All number divide by 8 also divide by 2 is 13

Ans. (b)

152. Sum of 1st n odd number

© The Institute of Chartered Accountants of India

S = 1+3+5+ ... + (2n – 1)

Since $S = \frac{n}{2}[2a+(n-1)d]$

S = n/2 [2.1 + (n – 1)2]

= n (1+n – 1)

= n(n)

$S = n^2$

Ans. (a)

153. Let the number be x.

According to the given condition of the problem is

36x = x + 1050

36x = 1050

$x = \frac{1050}{35}$

x = 30

Ans. (b)

154. The formula is $1^3+2^3+3^3+\ldots\ldots+n^3=\left\{\frac{n(n+1)}{2}\right\}^2$

$\therefore \quad 1^3+2^3+3^3+\ldots\ldots+12^3=\left\{\frac{12(12+1)}{2}\right\}^2$

$= \quad [6(13)]^2$

$= \quad [78]^2 = 6084$

Ans. (c)

155. Let $S = 1+9+24+46+75+\ldots\ldots+t_n$

shifting 1 places to the right in the RHS

$S = 1+9+24+46+75+\ldots\ldots\ldots t_{n-1}+t_n$

subtracting term by term,

$0=1+8+15+22+29+\ldots\ldots\ldots+(t_n-t_{n-1}) - t_n$

Transposing

t_n = 1+8+15+22+29+ ..... to n^{th} term

© The Institute of Chartered Accountants of India

$= \frac{n}{2}\{2.1+(n-1)7\}$

$= \frac{7}{2}n^2 - \frac{5}{2}n$

Now $Sn = \sum tn = \frac{7}{2}\sum n^2 - \frac{5}{2}\sum n$

$= \frac{7}{2}\frac{n(n+1)(2n+1)}{6} - \frac{5}{2}\frac{n(n+1)}{2}$

$= \frac{n(n+1)}{2}\left\{\frac{7}{6}(2n+1) - \frac{5}{2}\right\}$

$= \frac{n(n+1)}{2}\left(\frac{7}{3}n - \frac{4}{3}\right)$

$Sn = \frac{n(n+1)(7n-4)}{6}$

156. Let the number to be added be x then according to the problem.

$\frac{83+x}{263+x} = \frac{1}{3}$,

$3(83+x) = 263 + x$

$249 + 3x = 263 + x$

$3x - x = 263 - 249$

$2x = 14$

$x = \frac{14}{2} = 7$

Hence the required number is 7

Ans. (c)

157. Initially, let the number of employees be 9 and wages per head be Rs. 14. Then, total wages bill = Rs. (9×14) = Rs. 126

Further, the number of employees becomes 8 and the wages per head becomes Rs. 15.

∴ Now total wages bill = Rs. (8×15) = Rs. 120

∴ Ratio of the wages bill = 126:120

= 21:20

Thus the wages bill is decreased in the ratio 21:20

Ans. (c)

© The Institute of Chartered Accountants of India

ANSWERS

158. Let C gets Rs. = x

Given B = ¼ of C = ¼ (x)

and $A = \frac{2}{3}$ of B = 2/3 (¼x) = $\frac{1}{6}x$

Also, given $\frac{1}{6}x + \frac{1}{4}x + x = 680$

$$\frac{2x + 3x + 12x}{12} = 680$$

$$\frac{17x}{12} = 680$$

$$x = \frac{680 \times 12}{17} = 480$$

Ans. (c)

159. Let us assume that when x is added to each of the four given numbers, they become in proportion.

$\Rightarrow$ 10 + x : 18 + x = 22 + x : 38 + x

$\therefore$ Product of the means = Product of the extremes.

$\therefore$ (10+x) (38+x) = (18+x) (22+x)

$\Rightarrow$ $380+48x + x^2 = 396 + 40x + x^2$

8x = 16

x = 2

Required number = 2

Ans. (a)

160. Let the required numbers be x and y. Since the mean proportional between a and c is given by the relation $b = \sqrt{ac}$

$\therefore$ Mean proportional = $\sqrt{xy}$

According to the question,

$\sqrt{xy} = 24$

xy = 576 $\rightarrow$ (1)

Again suppose that the third proportional to x and y is z. Then

x : y = y : z

© The Institute of Chartered Accountants of India

$\Rightarrow$ x : z = y : y

$xz = y^2$

$\Rightarrow \quad z = \frac{y^2}{x}$

According to the question,

$\frac{y^2}{x} = 192$

$\Rightarrow \quad y^2 = 192\,x \qquad \rightarrow (2)$

From equation (2), $x = \frac{y^2}{192}$

Putting this value of x in equation (1)

$\frac{y^2}{192} . y = 576$

$y^3 = 576 \times 192$

$= \quad 24 \times 24 \times 24 \times 8$

$= \quad 24 \times 24 \times 24 \times 2 \times 2 \times 2$

$y = \quad (24 \times 24 \times 24 \times 2 \times 2 \times 2 \times 2)^{1/3}$

$= \quad 24 \times 2$

$y = 48$

(1) $\Rightarrow xy = 576$

$x\,(48) = 576$

$x = \frac{576}{48} = 8$

Hence the required number are 12 and 48.

161. $\lim\limits_{x \to 0} \frac{e^x + e^{-x} - 2}{x} \Rightarrow \left[\frac{e^{x-1}}{x} + \frac{(e^{-x} - 1}{x}\right]$

$\Rightarrow \lim\limits_{x \to 0} \left[\left(\frac{e^x - 1}{x}\right) + \left(\frac{e^{-x} - 1}{x}\right)\right] \qquad \left\{\lim\limits_{x \to 0} \frac{e^x - 1}{x} = 1\right\}$

$1 - 1 = 0$

Ans. (b) 0

© The Institute of Chartered Accountants of India

ANSWERS

162. $\underset{x\to 0}{\text{Lt}} \dfrac{e^{x^{-1}}-1}{e^{x^{-1}}+1} \Rightarrow \underset{x\to 0}{\text{Lt}} \left(\dfrac{e^{1/x}-1}{e^{1/x}+1}\right)$

RHL $\underset{x\to 0^+}{\text{Lt}} \left(\dfrac{e^{1/x}-1}{e^{1/x}+1}\right)$ App Lt = $\dfrac{\infty}{\infty}$

$\rightarrow$ RHL does not exist

LHL $\underset{x\to 0^-}{\text{Lt}} \left(\dfrac{e^{1/x}-1}{e^{1/x}+1}\right)$ App Lt $\dfrac{\infty}{\infty}$

LHL does not exist.

$\rightarrow$ Lt does not exist in f(x)

Ans. (c) does not exist

163. $\underset{x\to 0}{\text{Lt}} \dfrac{3x-|x|}{7x-5|x|}$

RHL $\underset{x\to 0^+}{\text{Lt}} \dfrac{3x-x}{7x-5x} = \dfrac{2x}{2x} = 1$

LHL $\underset{x\to 0^-}{\text{Lt}} \dfrac{3x-(-x)}{7x-5(-x)} = \dfrac{4x}{12x} = \dfrac{1}{3}$

LHL $\neq$ RHS

$\therefore$ f(x) does not exist at x = 0

Ans. (c) does not exist

164. $\underset{x\to 0}{\text{Lt}} \dfrac{e^{ax}-e^{bx}}{x} \Rightarrow \underset{x\to 0}{\text{Lt}} \left[a.\dfrac{\left(e^{ax}-1\right)}{ax} - b\dfrac{\left(e^{bx}-1\right)}{bx}\right]$

$\Rightarrow$ a.1 – b.1 $\Rightarrow$ a – b $\qquad \left\{\underset{x\to 0}{\text{Lt}} \dfrac{e^{x}-1}{x} = 1\right\}$

Ans. (a) a ... b

165. $\underset{x\to 0}{\text{Lt}} \dfrac{e^{x}-1}{\log(1+x)} \Rightarrow \underset{x\to 0}{\text{Lt}} \left[\left(\dfrac{e^{x}-1}{x}\right).\dfrac{x}{\log(1+x)}\right]$

$\Rightarrow \underset{x\to 0}{\text{Lt}} \left(\dfrac{e^{x}-1}{x}\right) . \underset{x\to 0}{\text{Lt}} \dfrac{1}{\dfrac{\log(1+x)}{x}}$

$\Rightarrow$ 1.1 = 1

Ans. (b) 1

© The Institute of Chartered Accountants of India

166. Given $y = \dfrac{\sqrt{1-x}}{\sqrt{1+x}}$

$$\frac{dy}{dx} = \frac{\left(\sqrt{1+x}\right)\frac{d}{dx}\left(\sqrt{1-x}\right) - \left(\sqrt{1-x}\right)\frac{d}{dx}\left(\sqrt{1+x}\right)}{\left(\sqrt{1+x}\right)^2}$$

$$= \frac{\left(\sqrt{1+x}\right)\left[\frac{1}{2}(1-x)^{\frac{1}{2}-1}(-1)\right] - \left(\sqrt{1-x}\right)\left[\frac{1}{2}(1+x)^{\frac{1}{2}-1}\right]}{(1+x)}$$

$$= \frac{-\frac{1}{2}\frac{\sqrt{1+x}}{\sqrt{1-x}} - \frac{1}{2}\frac{\sqrt{1-x}}{\sqrt{1+x}}}{(1+x)}$$

$$= -\frac{1}{2}\left[\frac{(1+x)+(1-x)}{\sqrt{1-x}\ \sqrt{1+x}} \cdot \frac{(1+x)}{1}\right]$$

$$\frac{dy}{dx} = -\frac{1}{2}\left[\frac{2}{\sqrt{1-x}\sqrt{1+x}} \cdot \frac{1+x}{1}\right] = \frac{-1}{(1+x)^{3/2}\left(\sqrt{1-x}\right)}$$

Ans. (b)

167. Given $y = \dfrac{x}{\sqrt{1+x^2}}$

$$\frac{dy}{dx} = \frac{\sqrt{1+x^2}(1) - x.\frac{1}{2\sqrt{1+x^2}}(2x)}{\left(\sqrt{1+x^2}\right)^2}$$

$$= \frac{\sqrt{1+x^2} - \frac{x^2}{\sqrt{1+x^2}}}{\left(1+x^2\right)}$$

$$= \frac{\left(1+x^2\right)-x^2}{\left(1+x^2\right)\left(\sqrt{1+x^2}\right)}$$

$$\frac{dy}{dx} = \frac{1}{\left(1+x^2\right)^{\frac{3}{2}}}$$

© The Institute of Chartered Accountants of India

$$\therefore \quad x^3 \frac{dy}{dx} = \frac{x^3}{\left(1+x^2\right)^{\frac{3}{2}}} = \left[\frac{x}{\sqrt{1+x^2}}\right]^3$$

$$x^3 \frac{dy}{dx} = [y]^3$$

Ans. (c)

168. Given $x^y = e^{x-4}$

Taking log on both sides

$\log \left(x^y\right) = \log \left(e^{x-y}\right)$

$y \log x = (x - y) \log e$

$y \log x = x - y$

$y (1 + \log x) = x$

$$y = \frac{x}{1+\log x}$$

Differentiate on both sides

$$\frac{dy}{dx} = \frac{(1+\log x).1 - x\left(\frac{1}{x}\right)}{(1+\log\ x)^2}$$

$$= \quad \frac{1+\log-1}{(1+\log x)^2} = \frac{\log x}{(1+\log x)^2}$$

Ans. (a)

169. Given $y^3 x^5 = (x+y)^8 \quad \rightarrow (1)$

Differentiate on both sides.

$$y^3\left(5x^4\right) + x^5 .3y^2 \frac{dy}{dx} = 8(x+y)^7\left[1+\frac{dy}{dx}\right]$$

$$5y^3x^4 + 3x^5y^2 \frac{dy}{dx} = 8(x+y)^7 + 8(x+y)^7 \frac{dy}{dx}$$

$$\frac{dy}{dx}\left[3x^5\ y^2 - 8\ (x+y)^7\right] = 8\ (x+y)^7 - 5y^3x^4$$

© The Institute of Chartered Accountants of India

$$\frac{dy}{dx} = \frac{8(x+y)^7 - 5y^3 x^4}{3x^5 y^2 - 8(x+y)^7} = \frac{8(x+y)^7 - 5\dfrac{(x+y)^8}{x}}{\dfrac{3}{y}(x+y)^8 - 8(x+y)^7}$$ Using equation (1)

$$= \frac{(x+y)^7\left[8 - \dfrac{5}{x}(x+y)\right]}{(x+y)^7\left(\dfrac{3}{y}(x+y) - 8\right)}$$

$$= \frac{y[8x - 5(x+y)]}{x[3(x+y) - 8y]} = \frac{y[8x - 5x - 5y]}{x[3x + 3y - 8y]}$$

$$= \frac{y[3x - 5y]}{x[3x - 5y]} = \frac{y}{x}$$

Ans. (a)

170. Given $y = x^{x^{x.....\infty}}$

i.e. $y = x^y$

Taking log on both sides

$\log y = \log \left(x^y\right)$

$\log y = y \log x$

Differentiate on both sides

$$\frac{1}{y}\frac{dy}{dx} = y.\frac{1}{x} + \log x.\frac{dy}{dx}$$

$$\frac{dy}{dx}\left[\frac{1}{y} - \log x\right] = \frac{y}{x}$$

$$\frac{dy}{dx}\left[\frac{1 - y \log x}{y}\right] = \frac{y}{x}$$

$$\frac{dy}{dx} = \frac{y^2}{x(1 - y \log x)}$$

$$\therefore \quad x.\frac{dy}{dx} = \frac{y^2}{1 - y \log x}$$

Ans. (b)

© The Institute of Chartered Accountants of India

ANSWERS

171. $\int_{-1}^{1}(e^x - e^x)dx = \int_{-1}^{1}(0)\,dx$

$= \quad 0$

Ans. (b) 0

172. $\int_{1}^{e}\frac{1+\log x}{x}\,dx$

Let $1 + \log x = t$ $\quad x = 0, t = 1$

$\frac{1}{x} = \frac{dt}{dx}$ $\quad x = e, t = 2$

$= \int_{1}^{2}+dt = \left[\frac{t^2}{2}\right]_1^2$

$= \quad 2 - \frac{1}{2} = \frac{3}{2}$

Ans. (a) $\frac{3}{2}$

173. $\int_{0}^{\log 3}\frac{ex}{1+e^x}dx$ $\quad$ If $\quad 1 + e^x = t$ $\quad x = 0, t = 2$

$e^x = \frac{dt}{dx}$ $\quad x = \log 3, t = 4$

$\int_{2}^{4}\frac{1}{-t}dt = [\log t]_2^4 \Rightarrow \log 4 - \log 2$

$= \log 2$

Ans. (b) log 2

174. $\int_{0}^{1}\frac{x}{1+\sqrt{1+x^2}}$ $\quad$ let $(1 + x^2) = t2$ $\quad x = 0, t = 1$

$x = 1\ t = \sqrt{2}$

$2x = 2t \quad \frac{dt}{dx} \Rightarrow dx = \frac{t}{x}\,dx$

$\int_{1}^{\sqrt{2}}\frac{t\,dt}{1+t} = \int_{1}^{\sqrt{2}}\left(1-\frac{1}{1+t}\right)dt$

© The Institute of Chartered Accountants of India

$= \quad \left[t - \log\left(1+t\right)\right]_1^{\sqrt{2}}$

$= \quad \left(\sqrt{2}-1\right) - \left[\log\left(1+\sqrt{2}\right) - \log 1\right]$

$= \quad \left(\sqrt{2}-1\right) - \log\left(1+\sqrt{2}\right) + 0$

$= \quad \left(\sqrt{2}-1\right) - \log\left(1+\sqrt{2}\right)$

Ans. (d) None of these

175. $\int_0^1 \frac{dx}{(1+x)(2+x)} \Rightarrow \int_0^1 \left(\frac{1}{1+x} - \frac{1}{2+x}\right) dx$

$\Rightarrow \left[\log(1+x) - \log(2+x)\right]_0^1 \Rightarrow \left[\log \frac{1+x}{2+x}\right]_0^1$

$\Rightarrow \log \frac{2}{3}$

Ans. (a) $\log \frac{2}{3}$

176. $a + ar = 15 \Rightarrow a(1+r) = 15$

$a = ar + ar^2 + ar^3 - \infty$

$a = \frac{ar}{1-r} \Rightarrow 1 - r = r \Rightarrow r = 1/2$

$\therefore a = \frac{15 \times 2}{3} = 10$

$\therefore$ Sum of Series $= \frac{a}{1-r} = \frac{10}{1-1/2} = 20$

Ans. (a) 20

177. $a = \frac{1}{1-x} \Rightarrow \frac{1}{a} = 1 - x$

$b = \frac{1}{1-y} \Rightarrow \frac{1}{b} = 1 - y$

$\therefore \frac{1}{a} + \frac{1}{b} = 1 - x + 1 - y \Rightarrow 2 - (x+y) = 2.1 = 1$

Ans. (c) 1

© The Institute of Chartered Accountants of India

ANSWERS

178. $90 = 2000 \times \frac{R}{100} \times \frac{3}{4}$

$\therefore$ R = 6%

Ans. (b) 6%

179. $216 = 5400 \times \frac{6}{100} \times n$

n = 4/6 yrs. = 8 months

Ans. (b) 8 months

180. $I_1 = 10000 \ \frac{R}{100} \times 2 = 200R$

$I_2 = 6000 \times \frac{R}{100} \times 3 = 1800R$

$I_1 + I_2 = 1900 \implies 200R + 180R = 1900$

$\therefore$ $380R = 1900$ $\therefore$ R = 5%

Ans. (b) 5%

181.If all the observation are equal.

Then standard deviation = 0

Ans. (a) 0

182. If every item is increased by 5 then mean ($\bar{x}$) also increased by 5, but the value of $\sum(x-\bar{x})^2$ remain same.

$\therefore$ Standard deviation will remain same,

Standard deviation = 10

Ans. (c) – 10

183. S.D. $= \sqrt{\frac{\sum d^2}{N}} = \sqrt{\frac{360}{10}} = 6$

Coefficient of variation $= 100 \ \frac{x\,SD}{Am}$

$= \frac{100 \times 6}{40} = 15$

Ans. (a) 15

© The Institute of Chartered Accountants of India

184. Coefficient of M.D. = $\frac{\text{M.D}}{\text{Am}} \times 100$

$44 = \frac{5.77 \times 100}{\text{AM}}$

A.M. = 13.11

Ans. 13.11

185. The S.D. of two values is equal to half their difference.

S.D. $= \frac{|a-b|}{2}$

The Statement is correct

Ans. (a) True

186. Computation of Correlation Coefficient

	x	y	xy	x^2	y^2
	50	40	2000	2500	1600
	50	40	2000	2500	1600
Total	100	80	4000	5000	3200

$\bar{x} = \frac{100}{2} = 50, \quad \bar{y} = \frac{80}{2} = 40$

Cov (x, y) $= \frac{4000}{2} - (50)(40) = 0$

$\therefore \quad r = 0$

Ans. (c)

187. Given $r_R = \frac{2}{3}, \quad \sum di^2 = 55$

$\therefore \quad r_R = 1 - \frac{b\sum di^2}{n\left(n^2-1\right)}$

$\frac{2}{3} = 1 - \frac{6(55)}{n\left(n^2-1\right)}$

$\frac{2}{3} - 1 = \frac{-330}{n\left(n^2-1\right)}$

$-\frac{1}{3} = -\frac{330}{n\left(n^2-1\right)}$

© The Institute of Chartered Accountants of India

$$n\left(n^2-1\right) = 990 = 10 \quad \left(10^2-1\right)$$

$\therefore$ n = 10 as n must a positive

Ans. (a)

188. Let us assume that $4x + 3y + 7 = 0 \quad \rightarrow (1)$

represent the regression line of x on y and $3x + 4y + 8 = 0 \quad \rightarrow (2)$ represent the regression line of y on x.

(1) $4x = -7 - 3y$

$$x = -\frac{7}{4} - \frac{3}{4}y$$

$$\therefore \text{ bxy} = -\frac{3}{4}$$

(2) $4y = -8 - 3x$

$$y = -2 - \frac{3}{4}x$$

$$\therefore \quad \text{byx} = -\frac{3}{4}$$

$$\therefore \quad r^2 = \text{byx. bxy} = \left(-\frac{3}{4}\right)\left(-\frac{3}{4}\right) = \frac{9}{16}$$

$$\therefore \quad r = \sqrt{\frac{9}{16}} = \pm\frac{3}{4} = -\frac{3}{4} = -0.75$$

(We take the sign of n as negative since both the regression coefficient are negative).

Ans. (c)

189. Ans. (d) Refer Properties

190. Given byx = 1.2 $\rightarrow (1)$

$$U = \frac{x-100}{2}$$

$\Rightarrow \quad x = 100 + 2U$

$\Rightarrow \quad \bar{x} = 100 + 2\bar{U}$

and $v = \dfrac{y-200}{3}$

$\Rightarrow y = 200 + 3v$

$\Rightarrow \bar{y} = 200 + 3\bar{v}$

© The Institute of Chartered Accountants of India

$$byx = \frac{\sum (x-\bar{x})(y-\bar{y})}{E(x-\bar{x})^2}$$

$$= \frac{\sum \left[2\left(U-\bar{U}\right).3\left(v-\bar{v}\right)\right]}{\sum \left[2(U-\bar{U})\right]^2}$$

$$= \frac{2\times 3}{4}\frac{\sum \left(U-\bar{U}\right)\left(V-\bar{V}\right)}{\sum (U-\bar{U})^2}$$

$= 3/2\ bvu$

$\Rightarrow$ bvu = 2/3 byx = 2/3 × 1.2 = 0.8

Ans. (b)

191. Let A, is First bag is selected

A2 is second bag is selected.

B: In a draw of 2 balls, one is red and the other is black.

The required probability

$P = P(A_1 \cap B) + P(A_2 \cap B)$

$= P(A_1)\, P(B/A_1) + P(A_2).\, P(B/A_2)$

Since there are two bags, the selection of each being equally likely.

$\therefore \quad P(A_1) = P(A_2) = 1/2$

$P(B/A_1)$ = Probability of drawing one red and one black ball in a draw of 2 balls from the 1st bag

$$= \frac{5C_1 \times 3C_1}{8C_2} = \frac{15}{28}$$

$P(B/A_2)$ = Probability of drawing one red and one black ball in a draw of 2 balls from the 2nd bags.

$$= \frac{4C_1 \times 5C_1}{9C_2} = \frac{5}{9}$$

(1) $\Rightarrow p = 1/2 \times \frac{15}{28} + \frac{1}{2} \times \frac{5}{9}$

$$p. = \frac{15}{56} + \frac{5}{18} = \frac{135+140}{504} = \frac{275}{504}$$

Ans. (a)

© The Institute of Chartered Accountants of India

ANSWERS

192. Let A1 is first purse is selected.

A2 is second purse is selected.

Let B: In a draw of one coin, one coin must be silver

The required probabilities.

$P = P\ (A_1 \cap B) + P\ (A_2 \cap B)$

$= \quad P(A_1).P(B/A_1) + P(A_2).P(B/A_2) \rightarrow (1)$

Since there are two purse, the selection of each being equally likely

$\therefore \quad P(A_1) = \frac{1}{2}, P(A_2) = \frac{1}{2}$

$P(B/A_1)$ = Probability of drawing one silver coin from the first purse.

$= \quad \frac{3C_1}{7C_1} = 3/7$

$P(B/A_2)$ = Probability of drawing one silver coin from the second purse.

$= \quad \frac{4C_1}{7C_1} = \frac{4}{7}$

Substituting $(1) \Rightarrow p=1/2 \left(\frac{3}{7}\right) + \frac{1}{2}\left(\frac{4}{7}\right)$

$= \quad \frac{3}{14} + \frac{4}{14} = \frac{7}{14} = \frac{1}{2}$

Ans. (a)

193. When two tosses of unbiased dice the total sample space.

S = {(1,1), (1,2), (1,3), (1,4), (1,5), (1,6)
(2,1), (2,2), (2,3), (2,4), (2,5), (2,6)
(3,1), (3,2), (3,3), (3,4), (3,5), (3,6)
(4,1), (4,2), (4,3), (4,4), (4,5), (4,6)
(5,1), (5,2), (5,3), (5,4), (5,5), (5,6)
(6,1), (6,2), (6,3), (6,4), (6,5), (6,6)}

n(s) = 36

In the above sample space, let x be the number of sines getting from the experiment.

Let x = 0, means no sin. = number of times = 25

x = 1, means no sin. = number of times = 10

© The Institute of Chartered Accountants of India

x = 2, means no sin. = number of times = $\frac{01}{36}$

∴ Expected table is:

x	0	1	2
p(x)	25	10	1

∴ The required probability = Mean = Expected Value

$= \sum \frac{x\, p(x)}{n(s)}$

$= \frac{0\times 25 + 1\times 10 + 2\times 1}{36}$

$= \frac{12}{36} = \frac{1}{3}$

Ans. (a)

194. The experiment of throwing three dice is theoretically same as that of throwing a die thrice.

Let E be the event of throwing six in a throw of die.

∴ P(E) = 1/6 and P ($\bar{E}$) = 1 – P(E)

= 1–1/6= 5/6

Let x denotes the random variable "number of Sixes".

∴ The possible values of x are 0, 1, 2, 3

∴ P(x=2) = P($E_1E_2\bar{E}_3$ or $E_1\bar{E}_2E_3$ or $\bar{E}_1E_2E_3$)

= $P(E_1).P(E_2)P(\bar{E}_3) + P(E_1)P(\bar{E}_2)P(E_3) + P(\bar{E}_1) + P(E_2)P(E_3)$

$= \frac{1}{6}.\frac{1}{6}.\frac{5}{6} + \frac{1}{6}.\frac{5}{6}.\frac{1}{6} + \frac{5}{6}.\frac{1}{6}.\frac{1}{6}$

$P(x=2) = \frac{15}{216}$

Ans. (c)

195. let A and B denote the events that the Chartered Accountant is selected in firms X and Y respectively. Then in the usual notations, we are given.

P(A) = 0.7

P($\bar{A}$) = 1– P(A) = 1=0.7=0.3

P($\bar{B}$) = 0.5

© The Institute of Chartered Accountants of India

$\therefore$ $P(B) = 1 - P(B) = 1 - 0.5 = 0.5$

and $P\left(\overline{A} \cup \overline{B}\right) = 0.6$

By De – Morgan's law

$\overline{(A \cap B)} = \overline{A} \cup \overline{B}$

$\therefore P\left(A \cup B\right) = 1 - P\,\overline{(A \cap B)}$

$= 1 - P\left(\overline{A} \cup \overline{B}\right)$

$= 1 - 0.6$

$= 0.4$

The probability that the Chartered Accountant will be selected in one of the two firms X or Y is given by:

$P(AUB) = P(A) + P(B) - P\left(A \cap B\right)$

$= 0.7 + 0.5 - 0.4$

$= 0.8$

Ans. (a)

196. $\int\left[\log(\log x) + \frac{1}{(\log x)^2}\right]dx$

$I = \log(\log x)\int dx - \int\left[\frac{d}{dx}\left[\log(\log x)\right]\right].x\,dx + \int\frac{1}{(\log x)^2} = dx$

$= \log(\log x).x - \int\frac{1}{(\log x)}.dx + \int\frac{1}{(\log x)^2}\,dx$

$x.\log(\log x) - \left[\frac{1}{(\log x)}\int dx + \left(\frac{1}{(\log x)^2}.\frac{x}{x}dx\right)\right] + \int\frac{1}{(\log x)^2} = dx$

$\Rightarrow x\log(\log x) - \frac{x}{\log x} - \int\frac{1}{(\log x)^2} + \int\frac{1}{(\log x)^2}dx$

$\Rightarrow x.\log(\log x) - \frac{x}{\log x} + c$

Ans. (a) $x.\log(\log x) - \frac{x}{\log x} + c$

197. 'is equal to' Satisfies Reflexive, Symnetric and transitive Relation

$\therefore$ This is Equivalence Relation.

Ans. (d) Equivalence Relation.

© The Institute of Chartered Accountants of India

198. $f(x) = x^2+2$

$\therefore\ f(-x) = (-x)^2 + 2$

$= \quad x^2 + 2$

$f(-x) = f(x)$

$\therefore$ f(x) is even function

Ans. (b) even function.

199. $f(x) = 12^{1+x} = 12.12^{x} \qquad 0 \leq x < 9$

Range = $12\times12°, \quad 12\times12^{1}$ .....12×12^{9}

$\therefore$ Range = $12 \leq f(x) < 12^{10}$

Ans. (a) $12 \leq f(x) \leq 12^{10}$

200. 'Is greater than' over the set of real number is not satisfied Reflexive and Symmetric relation it only satisfied transitive Relation

Ans. (a) Transitive relation.

Model Test Paper – BOS/CPT – 9

151. Given $\dfrac{x}{x+y} = \dfrac{17}{23}$

$\Rightarrow 23x = 17x + 17y$

$23x - 17x = 17y$

$6x = 17y$

$x = \dfrac{17}{6}y$

Now, $\dfrac{x+y}{x-y} = \dfrac{\frac{17}{6}y + y}{\frac{17}{6}y - y}$

$= \quad \dfrac{17y+6y}{6} \times \dfrac{6}{17y-6y}$

© The Institute of Chartered Accountants of India

$= \quad \frac{23y}{11y} = \frac{23}{11}$

$\therefore \quad \frac{x+y}{x-y} = \frac{23}{11}$

152. Given $\sqrt{1+\frac{25}{144}} = 1+\frac{x}{12}$

Squaring on both sides:

$1+\frac{25}{144} = \left(1+\frac{x}{12}\right)^2$

$\frac{144+25}{144} = 1+\frac{x^2}{144}+\frac{2x}{12}$

$\frac{169}{144} = \frac{144+x^2+24x}{144}$

$\therefore x^2+24x-25=0$

$(x+25)(x-1)=0$

$x=-25,\ x=1$

x = 1 ($\therefore$ negative neglected)

Ans. (a)

153. Given $(4)^3 \times \left(\sqrt{2}\right)^8 = 2^n$

i.e. $\left((2)^2\right)^3 \times \left((2)^{\frac{1}{2}}\right)^8 = 2^n$

$2^6 \times 2^4 = 2^n$

$2^{10} = 2^n$

$\therefore$ n = 10

Ans. (a)

154. Let total number of men went to a hotel = x

Given, A man Spent Rupees = Total number of men

$= \quad x$

$\therefore$ Given Data = x + x = 15625

© The Institute of Chartered Accountants of India

$x^2 = 15625$

$x = \sqrt{15625} = 125$

Ans. (b)

155. Given $A + B + C = 1000 \rightarrow (1)$

$A + C = 400 \rightarrow (2)$

$B + C = 700 \rightarrow (3)$

$(2) \Rightarrow A = 400 - C$

$(3) \Rightarrow B = 700 - C$

$(1) \Rightarrow 400 - C + 700 - C + C = 1000$

$C = 100$

Ans. (a)

156. Given $\log \left(\frac{a-b}{2}\right) = 1/2 (\log a + \log b)$

$\therefore \quad 2 \log \frac{a-b}{2} = \log a + \log b$

$\log \left(\frac{a-b}{2}\right)^2 = \log ab$

$\Rightarrow \quad \left(\frac{a-b}{2}\right)^2 = ab$

$\left(\frac{a-b}{4}\right)^2 = ab$

$(a - b)^2 = 4ab$

$a^2 + b^2 - 2ab = 4ab$

$a^2 + b^2 = 6ab$

Ans. (a)

157. Given $\log_{10} x = 4$

$\therefore x = 10^4$

Ans. (c) $x = 10000$

158. $\log 225 = \log (9 \times 25)$

$= \log 9 + \log 25$

© The Institute of Chartered Accountants of India

= $\log 3^2 + \log 5^2$

= $2\log 3 + 2 \log 5$

= $2\log 3 + 2 \log 10/2$

= $2\log 3 + 2 \log 10 - 2 \log 2$

= $2 \times 0.477 + 2 - 2\,(0.301)$

$\log 225 = 2.352$

Ans. (a)

159. Let $2^{100} = x$

Taking log on both sides.

$\log 2^{100} = \log x$

$100 \log 2 = \log x$

$\log x = 100 \times 0.3010$

$\log x = 30.1000$

$\therefore$ the no. of digits in 2^{100} is 31

Ans. (b)

160. Given $nP3 = \dfrac{n!}{(n-3)!} = 60$

$\therefore$ $n(n-1)(n-2) = 60 = 5 \times 4 \times 3$

$\therefore$ $n = 5$

Ans. (c)

161. $\underset{x\to 0}{\text{Lt}} \dfrac{a^x + b^x - 2}{x} \Rightarrow \underset{x\to 0}{\text{Lt}} \dfrac{(a^x - 1) + (b^x - 1)}{x}$

$\Rightarrow \underset{x\to 0}{\text{Lt}} \dfrac{a^x - 1}{x} + \underset{x\to 0}{\text{Lt}} \dfrac{b^x - 1}{x} \qquad \left[\underset{x\to 0}{\text{Lt}} \dfrac{a^x - 1}{x} = \log_e^a\right]$

App Lt

= $\log a + \log b = \log (ab)$

Ans. (a) $\log (ab)$

162. $\underset{x\to 0}{\text{Lt}} \dfrac{10^x - 5^x - 2^x + 1}{x}$

$\Rightarrow \underset{x\to 0}{\text{Lt}} \left[\dfrac{(10^x - 1)}{x} - \dfrac{(5^x - 1)}{x} - \dfrac{(2^x - 1)}{x}\right] \qquad \left\{\underset{x\to 0}{\text{Lt}} \dfrac{a^x - 1}{x} = \log_e^a\right\}$

© The Institute of Chartered Accountants of India

App Lt

= $\log 10 - \log 5 - \log 2$

= $\log\left(\frac{10}{5\times 2}\right) = \log 1 \Rightarrow 0$

Ans. (b) 0

163. $\underset{x\to 0}{\text{Lt}} \frac{10^x - 5^x - 2^x - 1}{x^2} \Rightarrow \underset{x\to 0}{\text{Lt}} \frac{[5^x \times 2^x - 5^x - 2^x - 1]}{x^2} = \frac{5^x(2^x-1) - 1(2^x-1)}{x^2}$

$\underset{x\to 0}{\text{Lt}} \frac{(2^x-1)(5^x-1)}{x^2} \Rightarrow \underset{x\to 0}{\text{Lt}} \frac{(2^x-1)}{x} \times \underset{x\to 0}{\text{Lt}} \frac{(5^x-1)}{x}$

$\Rightarrow \log 2 \times \log 5$

Ans. (a) $\log 5 \times \log 2$

164. $\underset{x\to 0}{\text{Lt}} \frac{e^{5x} - e^{3x} - e^{2x} + 1}{x}$

$\Rightarrow \underset{x\to 0}{\text{Lt}} \left[5\frac{(e^{5x}-1)}{5x} - 3.\frac{(e^{3x}-1)}{3^x} - 2\frac{(e^{2x}-1)}{2^x}\right]$

App Lt

5.1 – 3.1 – 2.1

= 5 – 3 – 2 = 0 $\qquad \left\{\underset{x\to 0}{\text{Lt}} \frac{e^x - 1}{x} = 1\right\}$

Ans. (b) 0

165. $\underset{x\to 0}{\text{Lt}} \frac{e^{5x} - e^{3x} - e^{2x} - 1}{x^2} = \underset{x\to 0}{\text{Lt}} \frac{(e^{3x}.e^{2x} - e^{3x} - e^{2x} - 1)}{x^2}$

$\underset{x\to 0}{\text{Lt}} \frac{\left[e^{3x}(e^{2x}-1) - 1(e^{2x}-1)\right]}{x^2} \Rightarrow \underset{x\to 0}{\text{Lt}} \frac{\left[(e^{3x}-1)\,(e^{2x}-1)\right]}{x^2}$

$\underset{x\to 0}{\text{Lt}} \left(\frac{e^{3x}-1}{x}\right) \times \underset{x\to 0}{\text{Lt}} \left(\frac{e^{2x}-1}{x}\right)$

App Lt

$3 \times 2 = 6$

Ans. (a) 6

© The Institute of Chartered Accountants of India

ANSWERS

166. Given $\int \frac{1}{\sqrt{x^2+a^2}}\,dx$

Let $\sqrt{x^2+a^2} = z - x$

$\therefore \quad z = x + \sqrt{x^2+a^2}$

$\frac{dz}{dx} = 1 + \frac{1}{2\sqrt{x^2+a^2}}(2x)\,dx$

$= \quad \frac{\sqrt{x^2+a^2}+x}{\sqrt{x^2}+a^2} = \frac{z}{\sqrt{x^2+a^2}}$

$\therefore \quad \frac{dz}{z} = \frac{dx}{\sqrt{x^2+a^2}}$

$\therefore \quad \int \frac{dx}{\sqrt{x^2+a^2}} = \int \frac{dz}{z} = \log z + c$

$= \quad \log\left(x+\sqrt{x^2+a^2}\right) + c$

167. $\int \frac{1}{\sqrt{x^2-a^2}}dx$

Let $\sqrt{x^2-a^2} = z - x$

$\therefore \quad z = x + \sqrt{x^2-a^2}$

$\frac{dz}{dx} = 1 + \frac{1}{2\sqrt{x^2-a^2}}(2x) = 1 + \frac{x}{\sqrt{x^2-a^2}}$

$= \quad \frac{\sqrt{x^2-a^2}+x}{\sqrt{x^2-a^2}} = \frac{z}{\sqrt{x^2-a^2}}$

$\frac{dz}{z} = \frac{dx}{\sqrt{x^2-a^2}}$

$\therefore \quad \int \frac{1}{\sqrt{x^2-a^2}}\,dx = \int \frac{1}{2}dz + c$

$= \quad \log(z) = \log\left(x+\sqrt{x^2-a^2}\right) + c$

Ans. (c)

© The Institute of Chartered Accountants of India

168. Let $t = 3x$

$\therefore\ dt = 3\ dx$

$\therefore\ I = \int \frac{1}{t^2 - 1}\frac{dt}{3} = \frac{1}{3}\int \frac{1}{t^2 - 1^2}\,dt$

$= \frac{1}{3}\frac{1}{2+1}\log\left(\frac{t-1}{t+1}\right) + c$

$= \frac{1}{6}\log\left(\frac{3x-1}{3x+1}\right) + c$

Ans. (b)

169. Let $I = \int \frac{x-1}{\sqrt{x^2+1}}\,dx$

$\therefore\ I = \int\left[\frac{x}{\sqrt{x^2+1}} - \frac{1}{\sqrt{x^2+1}}\right]dx$

$= \int \frac{x}{\sqrt{x^2+1}}\,dx - \int \frac{1}{\sqrt{x^2+1}}\,dx$

$I = I_1 - I_2$ (say)

$I_1 = \int \frac{x}{\sqrt{x^2+1}}\,dx$

Let $t = x^2 + 1$

$\therefore\ dt = 2x\ dx$

$\therefore\ I_1 = \int \frac{1}{\sqrt{t}}\frac{dt}{2}$

$= \frac{1}{2}\int t^{-1/2}\,dt$

$= \frac{1}{2}\frac{t^{1/2}}{1/2} = \sqrt{t} = \sqrt{x^2+1}$

$I_2 = \int \frac{1}{\sqrt{x^2+1}}\,dx$

$= \log\left(x + \sqrt{x^2+1}\right)$

© The Institute of Chartered Accountants of India

$\therefore \quad I = I_1 - I_2$

$I = \sqrt{x^2+1} - \log\left(x+\sqrt{x^2+1}\right) + c$

Ans. (a)

170. Let $I = \int (1-x^2)\log x \, dx$

$\therefore I = \int \log x \,(1-x^2)\, dx$

[Here (log x) i s to be take n as first function and $(1 - x^2)$ as second function]

Integrating by parts:

$I = \log x \left(x - \frac{x^3}{3}\right) - \int \frac{1}{x}\left(x - \frac{x^3}{3}\right) dx$

$= \left(x - \frac{x^2}{3}\right) x \log x - \int \left(x - \frac{x^2}{3}\right) dx$

$= \left(1 - \frac{x^2}{3}\right) x \log x - \int \left(x - \frac{x^3}{3*3}\right) + c$

$I = \left(1 - \frac{x^2}{3}\right) x \log x - \left(x - \frac{x^3}{9}\right) + c$

Ans. (c)

171. Among 4 doctors, 4 officers and 1 doctor who is also an officer committee of 3 can be form in such manner.

(i) 1 doctor, 1 officer, 1 doctor who is also officer $= 4_{C_1} \times 4_{c_1} \times 1 = 16$

(ii) 2 doctor and doctor – officer = $4_{C_2} \times 1 = 6$

(iii) 2 officer and doctor – officer = ${}^4C_2 \times 1 = 6$

(iv) 2 doctor and 1 officer = $4_{C_2} \times 4_{C_1} = 24$

(v) 1 doctor and 2 officer = $4_{C_1} \times 4_{C_2} = 24$

Total no. of ways = 16 + 6 + 6 + 24 + 24 = 76

Ans. (a) 76

172. Elector can vote for one or more vacancies in such manner …

(i) For 3 vacancies – $5_{C_3} = 10$

© The Institute of Chartered Accountants of India

(ii) For 2 vacancies – $5_{C_2} = 10$

(iii) For 1 vacancy – $5_{C_1} = 5$

$\therefore$ Total ways = 10 + 10 + 5 = 25

Ans. (c) 25

173. No. of ways in which 12 different thing distributed in 4 groups.

$= \dfrac{12!}{(3!)^4} = 15400$

Ans. (a) 15,400

174. Factor of 420 is = {2, 3, 4, 5, 6, 7, 10, 12, 14, 15, 20, 21, 30, 28, 35, 42, 60, 84, 105, 210, 140, 420}

No. of factor of 420 = 22

Ans. (b) 22.

175. Five balls are kept in 3 boxes as no box will empty

$= \left(5_{C_1} \times 4_{c_1} \times 3_{c_3}\right) \times 3!$

$= (5 \times 4 \times 1) \times 6 = 120$ ways.

Ans. (b) 120 ways.

176. 243 + 324 + 432 + – n terms

$3^5.1 + 3^4.4 + 3^3.4^2 +$ – n terms

$\therefore a = 3^5 \qquad r = 4/3$

$$Sn = \frac{3^5.\left[\left(\frac{4}{3}\right)^n - 1\right]}{\left(\frac{4}{3} - 1\right)} = 3^5.3\left[\left(\frac{4}{3}\right)^n - 1\right]$$

$$= 3^6\left[\frac{4^n}{3^n} - 1\right]$$

Ans. (a) $3^6\left[\dfrac{4^n}{3^n} - 1\right]$

177. $S_8 = 5.S_4 \Rightarrow \dfrac{a\left[r^8 - 1\right]}{r - 1} = \dfrac{5.a\left[r^4 - 1\right]}{r - 1}$

$\Rightarrow \left(r^4 + 1\right) = 5$

© The Institute of Chartered Accountants of India

$r^4 = 4 = \left(\sqrt{2}\right)^4$

$\therefore r = \pm \sqrt{2}$

Ans. (c) $\pm \sqrt{2}$

178. 4+44+444 ... n terms

$= \frac{4}{9}[9 + 99 + 999 + ...\text{n terms}]$

$= \frac{4}{9}[(10-1)+(100-1)+(1000-1)+ ... \text{ n terms})$

$= \frac{4}{9}\left[\frac{10(10^n-1)}{10-1} - n\right] \Rightarrow \frac{4}{9}\left[\frac{10(10^n-1)}{9} - n\right]$

Ans. (a) $\frac{4}{9}\left[\frac{10(10^n-1)}{9} - n\right]$

179. $\frac{a+b}{2} = 15$ and $\sqrt{ab} = 9 \therefore ab = 81$

$a = (30 - b)$ & $(30 - b)b = 01$

$\therefore b^2 - 30b + 81 = -0 \therefore b = 27, 3$ and $a = 3, 27$

$\therefore$ Nos are 27, 3

Ans. (a) 27, 3

180. Product of n Gm between two No. is equal to n th Power of single Gm between two nos. This statement is correct.

Ans. (a) True

181. The weighted arithmetic mean of first a natural numbers whose weights are equal to the corresponding number is equal to

$\frac{2n+1}{3}$

Ans. (a) $\frac{2n+1}{3}$

182. $\bar{x} = \frac{w_1x_1 + w_2x_2 + w_3x_3}{(x_1 + x_2 + x_3)}$

$110 = \frac{100 \times 5 = 125 \times 5 + w_3 \times 5}{15}$

© The Institute of Chartered Accountants of India

$1650 = 500 + 625 + 5.w_3$

$w_3 = \frac{525}{5} = 105$ kg.

Ans. (b) 105 Kgs.

183. $\sum x - n \times 2.5 = 50$

$\sum x - 2.5n = 50 \rightarrow$ (i)

$\sum x - 3.5n = -50 \rightarrow$ (ii)

[Eg. (i) – Eg. (ii)]

$1.0n = 100$

$\therefore n = 100 \qquad \therefore \Sigma x = 300$

$\therefore$ mean $= \frac{\Sigma x}{n} = \frac{300}{100} = 3$

Ans. (a) 100, 3

184. The most reliable value is mean.

Ans. (a) Mean

185. In which Central Value arranging is required – Median

Ans. (c) Median

186. There are 365 days in a normal year (without leap year)

No. $365 = 7 \times 52 + 1$

$\therefore$ In a year will contain at least 52 Tuesday

The possible remaining one Tuesday

Let A be the event of getting 53 Tuesday in the year.

$\therefore$ P (A) = 1/7

Ans. (b)

187. Given two unbiased dice are thrown, then the simple space are:

S = { (1,1), (1,2), (1,3), (1,4), (1,5), (1,6),
(2,1), (2,2), (2,3), (2,4), (2,5), (2,6),
(3,1), (3,2), (3,3), (3,4), (3,5), (3,6),
(4,1), (4,2), (4,3), (4,4), (4,5), (4,6),
(5,1), (5,2), (5,3), (5,4), (5,5), (5,6),
(6,1), (6,2), (6,3), (6,4), (6,5), (6,6)}

© The Institute of Chartered Accountants of India

$\therefore$ n(S) = 36

Sample space of sum of the faces is not less than 10

A = {(4,6), (5,5), (5,6), (6,4), (6,5), (6,6)}

n(A) = 6

$\therefore$ Required probability = $\frac{n(A)}{n(S)} = \frac{6}{36} = \frac{1}{6}$

Ans. (a)

188. Let A be the person travels by a plane

$\therefore$ P(A) = $\frac{1}{5}$

Let B be the person travels by a train

$\therefore$ P (B) = $\frac{2}{3}$

$\therefore$ Probability of his travelling neither by plane nor by train.

P(AB) = P(A). P(B) (Since A and B are mutually exclusive conditional probability)

= $\left(\frac{1}{5}\right)\left(\frac{2}{3}\right) = \frac{2}{15}$

Ans. (b)

189. Let A denote the event of drawing a diamond and B denote the event of drawing a King from a pack of Cards. Then we have P(A) = $\frac{13}{52} = \frac{1}{4}$

and P(B) = $\frac{4}{52} = \frac{1}{13}$

P(AUB) = P(A) + P(B) – P(A $\cap$ B)

= $\frac{1}{4} + \frac{1}{13} - P(A \cap B) \rightarrow (1)$

There is only one case favourable to the event A $\cap$ B vize, king of diamond.

Hence, P (A $\cap$ B) = $\frac{1}{52}$

$\therefore$ (1) $\Rightarrow$ P(AUB) = $\frac{1}{4} + \frac{1}{13} - \frac{1}{52} = \frac{13+4-1}{52} = \frac{16}{52} = \frac{4}{13}$

Ans. (c)

© The Institute of Chartered Accountants of India

190. Let us define the events:

E_1 : A solves the problem

E_2 : B solves the problem

then we are given

$$P(E_1) = \frac{6}{6+9} = \frac{6}{15} = \frac{2}{5}$$

and $P(E_2) = \frac{10}{10+12} = \frac{5}{11}$

Assuming that A and B try to solve the problem independently. E_1 and E_2 are independent.

$$\therefore P(E_1 \cap E_2) = P(E_1).P(E_2) = \frac{2}{5} \times \frac{5}{11} = \frac{2}{11}$$

The problem will be solved if at least one of the students A and B solves the problem.

Hence, the probability of the problem being solved is given by

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2)$$

$$= \frac{2}{5} + \frac{5}{11} - \frac{2}{11}$$

$$= \frac{22 + 25 - 10}{55} = \frac{37}{55} = 0.673$$

Ans. (a)

192. Given Mean of Binomial distribution = $\mu = np = 3 \rightarrow (1)$

and Variance of Binomial distribution = $\sigma^2 = npq = .2 \rightarrow (2)$

$$\frac{(2)}{(1)} \Rightarrow \frac{npq}{np} = \frac{2}{3}$$

$\therefore q = 2/3$

$\therefore p + q = 1$

$$p = 1 - q = 1 - \frac{2}{3} = \frac{1}{3}$$

$(1) \Rightarrow n(1/3) = 3$

$= n = 9$

$\therefore p=1/3, q = 2/3, n=9$

By the Binomial distribution $p(x) = nC_x\, p^x\, q^{n-x}$. The probability that the variate takes values less than or equal to 2

© The Institute of Chartered Accountants of India

i.e. $p(x \leq 2) = p(x = 0) + P(x = 1) + P(x = 2)$.

$= {}^{9}C_0 (1/3)^0 (2/3)^{9-0}$

$+ {}^{9}C_1 (1/3)^1 (2/3)^{9-1}$

$+ {}^{9}C_2 (1/3)^2 (2/3)^{9-2}$

$= {}^{9}C_0 (1) (2/3)^9$

$+ {}^{9}C_1 (1/3)^1 (2/3)^8 + {}^{9}C_2 (1/3)^2 (2/3)^7$

$= (2/3)^9 + 3 (2/3)^8 + 4 (2/3)^7$

$P(x \leq 2) = 0.3767$

Ans. (a)

193. Exhaustive cases: 2 digits can be selected out of 9 digits 1 through 9 in ${}^{9}C_2$ ways.

$\therefore$ Exhaustive number of cases $= {}^{9}C_2 = \frac{9 \times 8}{1 \times 2} = 36$

Favoarable number of cases. Among the digits 1 through 9.

Even digits are: 2, 4, 6 and 8 i.e. 4 in all

Odd digits are 1, 3, 5, 7 and 9 i.e. 5 in all.

The sum of the two digits drawn will be even if

(i) Either both the selected digits are even (or)

(ii) both the selected digits are odd.

Two even digits can be selected out of the 4 even digits in ${}^{4}C_2$ ways and two odd digits can be selected out of the 5 odd digits in ${}^{5}C_2$ ways.

Hence, the favourable number of cases that the sum of the two selected digits in even.

$= {}^{4}C_2 + {}^{5}C_2$

$= \frac{4 \times 3}{1 \times 2} + \frac{5 \times 4}{1 \times 2}$

$= 6+10 = 16$

$\therefore$ P (sum of the two selected digits is even)

$= \frac{\text{Number of favourable cases}}{\text{Exhaustive number of cases}}$

$= \frac{16}{36} = \frac{4}{9}$

and P(Both selected digits are odd) $= \frac{{}^{5}C_2}{{}^{9}C_2} = \frac{10}{36} = \frac{5}{18}$

Ans. (e)

© The Institute of Chartered Accountants of India

194. Let S be the sample space of the experiment

$\therefore$ S = {(1,1), (1,2), ... (6,5), (6,6)}

Let A = event of getting sum 6

and B = event of getting 4 at least once.

$\therefore$ A = {(1,5), (2,4), (3,3), (4,2), (5,1)}

and B = {(4,1), (4,2), (4,3), (4,4), (4,5), (4,6), (1,4), (2,4), (3,4), (5,4), (6,4)}

$\therefore$ P(A) = 5/36 and P(B) = $\frac{11}{36}$

Also, AB = {(4,2), (2,4)}

$\therefore$ P(AB) = $\frac{2}{36}$

$\therefore$ The required probability

= Probability of getting 4 on at least one die given that sum is 6

= $P(B|A) = \frac{P(BA)}{P(A)} = \frac{P(AB)}{P(A)}$

= $\frac{2/36}{5/36} = \frac{2}{5}$

Ans. (b)

196. Standard Error of mean = $\frac{\sigma}{\sqrt{n}}$

= $\frac{12.6}{\sqrt{36}} = \frac{12.6}{6}$

Standard Error of mean = 2.1

Ans. (a) 2.1

197. Standard Error of Mean without replacement

= $\frac{\sigma}{\sqrt{n}}\sqrt{\frac{N-n}{N-1}}$

= $\frac{12.6}{\sqrt{36}}\sqrt{\frac{101-36}{101-1}} = 2.1\times\sqrt{0.65}$

SE = 2.1 × 0.806 = 1.69

Ans. (b) 1.69

© The Institute of Chartered Accountants of India

ANSWERS

198. $P = \frac{5}{25} = \frac{1}{5}$

$Q = 1 - \frac{1}{5} = \frac{4}{5}$

$n = 5$

S.E. of proportion of defectives $= \sqrt{\frac{PQ}{n}}$

$= \sqrt{\frac{1}{5} \times \frac{4}{5} \times \frac{1}{5}} = \sqrt{0.032}$

SE = 0.1088

Ans. (b) 0.1088

199. n = 2, N = 4

Total number of possible sample of size of with replacement = $4^2 = 16$

Ans. (a) 16

200. n = 2, N = 4

Total number of possible sample of size without replacement = $N_{C_n} = 4_{C_2} = 6$

Ans. (b) 6

Model Test Paper – BOS/CPT – 10

151. The value of $3^3 + 4^3 + 5^3 + ... + 11^3$

$= \left[\frac{11(11+1)}{2}\right]^2 - \left[1^3 + 2^3\right]$

$= (11 \times 6)^2 - (1 + 8)$

$= (66)^2 - (9)$

$= 4356 - 9 = 4347$

Ans. (c)

152. Let the two numbers are x and y

Given $x + y = 75 \rightarrow (1)$

$x - y = 20 \rightarrow (2)$

© The Institute of Chartered Accountants of India

(1)+(2) $\Rightarrow 2x = 95$

$$x = \frac{95}{2}$$

$\therefore$ (1) $\Rightarrow = 75 - \frac{95}{2}$

$$= \frac{55}{2}$$

$\therefore$ The difference of their squares = $x^2 - y^2$

$$= \left(\frac{95}{2}\right)^2 - \left(\frac{55}{2}\right)^2$$

$$= \frac{9025}{4} - \frac{3025}{4}$$

$$= \frac{6000}{4} = 1500$$

Ans. (a)

153. Let the two numbers are x and y

Given $x + y = 13 \rightarrow (1)$

and $x^2 + y^2 = 85 \rightarrow (2)$

(1) $\Rightarrow$ $y = 13 - x$

Substitute $y = 13 - x$ in (2)

$x^2 + (13 - x)^2 = 85$

$x^2 + 169 + x^2 - 26x - 85 = 0$

$2x^2 - 26x + 84 = 0$

$x^2 - 13x + 42 = 0$

$(x - 7)(x - 6) = 0$

x =7, x = 6

When x = 7 (1) $\Rightarrow$ y = 13 – 7 = 6

When x = 6 (1) $\Rightarrow$ y = 13 – 6 = 7

$\therefore$ The number (7, 6)

Ans. (a)

© The Institute of Chartered Accountants of India

ANSWERS

154. Let the two consecutive members are x and x – 1

Given $x^2 - (x-1)^2 = 37$

$x^2 - (x^2 + 1 - 2x) = 37$

$x^2 - x^2 - 1 + 2x = 37$

$2x = 38$

$x = 19$

$\therefore x - 1 = 19 - 1 = 18$

Ans. (a)

155. Let the number be x.

Given condition (1) $\frac{x}{x+3} \rightarrow (1)$

Given condition (2) $\Rightarrow \frac{x+7}{x+3-2} = 2$

$x + 7 = 2(x+1)$

$= 2x + 2$

$x = 5$

$\therefore (1) \Rightarrow \frac{5}{5+3} = \frac{5}{8}$

156. Given $\log_x \sqrt{3} = \frac{1}{6}$

$x^{\frac{1}{6}} = \sqrt{3}$

$x = \left(\sqrt{3}\right)^6$

$= \left(\sqrt{3}\right)^2 \left(\sqrt{3}\right)^2 \left(\sqrt{3}\right)^2$

$= 3 \times 3 \times 3$

$x = 27$

Ans. (b)

157. Let $y = a^{\log a^x}$

Taking log on both sides

$\log y = \log \left[a^{\log a^x}\right]$

© The Institute of Chartered Accountants of India

= $\log_a x . \log a$

$\log y = \log x$ [$\because \log_b^a . \log_c^b = \log_c^a$ By properties]

Taking Exponent on both sides.

$e^{\log y} = e^{\log x}$

$y^{\log e} = x^{\log x}$

$y = x$

Ans. (a)

158. Let $y = 3^{2-\log_3 6}$

= $3^2 . 3^{-\log_3 6}$

= $9.3^{-[\log_3 (3x2)]}$

= $9.3^{-[\log_3 3+\log_3 2]}$

= $9.3^{-[1+\log_3 2]}$

= $9.3^{-1}.3^{-\log 3^2}$

= $3.3^{\log_3 (1/2)}$

= $3.(1/2)$ [$\because$ a $\log_a x$ = x properties]

$\therefore$ $y = 3/2$

Ans. (b)

159. $\log 30 = \log (2\times3\times5)$

= $\log 2 + \log 3 + \log 5$

= $0.3010 + 0.4771 + 0.6990$

= 1.4771

Ans. (c)

160. $\log_{10} 124.5 + \log_{10} 379 = \log_{10}(12.45 \times 10)$

+ $\log_{10}(3.79 \times 100)$

= $\log_{10} 12.45 + \log_{10} 10$

+ $\log_{10} 3.79 + \log_{10} 100$

= $1.0952 + 0.5786 + 2$

$\therefore$ $\log_{10} 124.5 + \log_{10} 379 = 4.6738$

Ans. (b)

© The Institute of Chartered Accountants of India

ANSWERS

161. $n_{P_5} : n_{P_3} = 2:1 \Rightarrow \frac{n!}{(n-5)!} \div \frac{n!}{(n-3)!} = \frac{2}{1}$

$\Rightarrow \quad \frac{(n-3)!}{(n-5)!} = \frac{2}{1} \Rightarrow (n-3)(n-4) = 2.1$

$\therefore \quad n - 3 = 2 \Rightarrow n = 5$

Ans. (b) 5

162. No. of ways to enter into room = 10

No. of ways to came out from a different door = 9

$\therefore$ Total Ways = 10.9 = 90 ways

Ans. (a) 90

163. Five digit Nos. by using digit (1, 2, 3, 4, 6) = 5! = 120

Four digit nos. by using digit (1, 2, 3, 4, 6) = $5 P_4$

= 120

$\therefore$ Total Nos. greater than 1000 = 120 + 120

= 240

Ans. (c) 240

164. Total Nos. of 6 digit (greater than 1 lakh)

by using (1, 1, 1, 2, 2, 3) are = $\frac{6!}{3!\ 2!}$

= 60

Ans. (a) 60

165. No. of ways in which 17 billiard can be arranged.

If 7 are black, 6 red and 4 white are

= $\frac{17!}{7!\ 6!\ 4!} = 4084080$

Ans. (b) 4084080

166. Let $I = \int \frac{xe^x}{(x+1)^2} dx$

$= \int \frac{(x+1-1)e^x}{(x+1)^2} dx$

© The Institute of Chartered Accountants of India

$$= \int \frac{(x+1)e^x}{(x+1)^2}dx - \int \frac{e^x}{(x+1)^2}dx$$

$$= \int \frac{e^x}{(x+1)}dx - \int \frac{e^x}{(x+1)^2}dx$$

$$= I_1 - I_2$$

$$I = \int \frac{e^x\,dx}{(x-1)}$$

Integrating by parts.

$$I_1 = \frac{1}{1+x}e^x - \int\left(-\frac{1}{(x+1)^2}\right)e^x\,dx$$

$$\therefore I = \frac{1}{(1+x)}e^x + \int \frac{1}{(x+1)^2}e^x\,dx - \int \frac{e^x}{(x+1)^2}dx$$

$$I = \frac{1}{(1+x)}e^x$$

Ans. (b)

167. $\int e^x \frac{(x-1)}{(x+1)^3}dx = \int \frac{x+1-2}{(x+1)^3}e^x\,dx$

$$= \int e^x \left\{\frac{1}{(x+1)^2} - \frac{2}{(x+1)^3}\right\}dx$$

$$= \int e^x \{f(x) + f'(x)\}dx$$

Where $f(x) = \frac{1}{(x+1)^2}$

$$= e^x f(x) + c$$

$$\int \frac{e^x(x-1)}{(x+1)^3}dx = \frac{ex}{(x+1)^2} + c$$

Ans. (c)

168. See the text book example page No. 9.28

Ans. (a)

© The Institute of Chartered Accountants of India

ANSWERS

169. $\int \frac{dx}{x^2 - a^2} = \int \frac{dx}{(x-a)(x+a)}$

$\int \frac{1}{2a}\left(\frac{1}{x-a} - \frac{1}{x+a}\right) dx$

$= \frac{1}{2a}\left[\int \frac{dx}{x-a} - \int \frac{dx}{x+a}\right]$

$= \frac{1}{2a}[\log(x-a) - \log(x+a)]$

$\int \frac{dy}{x^2 - a^2} = \frac{1}{2a} \log\left[\frac{x-a}{x+a}\right]$

170. $\int \frac{1}{a^2 - x^2} dx = \int \frac{dx}{(a+x)(a-x)}$

$= \int\left(\frac{-1}{2a}\right)\left[\frac{1}{a+x} - \frac{1}{a-x}\right] dx$

$= -\frac{1}{2a}\left[\int \frac{1}{a+x} dx - \int \frac{1}{a-x} dx\right]$

$= -\frac{1}{2a}[\log(a+x) - \log(a-x)]$

$= \frac{-1}{2a} \log\left[\frac{a+x}{a-x}\right]$

171. $e^{x-y} + \log xy + xy = 0$

d.wr.t.u.

$e^{x-y}\left(1 - \frac{dy}{dx}\right) + \frac{1}{xy}\left[x\frac{dy}{dx} + y.1\right] + \left[x\frac{dy}{dx} + y.1\right] = 0$

$e^{x-y} - e^{x-y}.\frac{dy}{dx} + \frac{1}{y}\frac{dy}{dx} + \frac{1}{x} + x\frac{dy}{dx} + y = 0$

© The Institute of Chartered Accountants of India

$$\left(-e^{x-y}+\frac{1}{y}+x\right)\frac{dy}{dx} = -e^{x-y}-\frac{1}{x}-y$$

$$\frac{1}{y}(xy+1-y.e^{x.y})\frac{dy}{dx} = -\frac{1}{x}(x.e^{x-y}+1+xy)$$

$$\therefore \frac{dy}{dx}=\frac{-y}{x}.\left(\frac{x.e^{x-y}+1+xy}{1+xy-y.e^{x-y}}\right)$$

Ans. (d) None of these

172. $y = x^{\log(\log x)}$

$\log y = \log (\log x) . \log x$

$$\frac{1}{y}.\frac{dy}{dx} = \log (\log x).\ \frac{1}{x} + (\log x).\ \frac{1}{(\log x)}.\frac{1}{x}$$

$$\therefore \quad \frac{dy}{dx}=\frac{y}{x}\ [\log (\log x) + 1]$$

Ans. (a) $\frac{y}{x}\ [\log (\log x) + 1]$

173. $y = x + \dfrac{1}{x+\dfrac{1}{x}}$

$$y = \frac{x^3+x+x}{x^2+1}=\frac{x^3+2x}{x^2+1}$$

$$\frac{dy}{dx}=\frac{\left(x^2+1\right)\left(3x^2+2\right)-\left(x^3+2x\right)(2x)}{\left(x^2+1\right)^2}$$

$$= \quad \frac{3x^4+3x^2+2x^2+2-2x^4-4x^2}{\left(x^2+1\right)^2}$$

$$\frac{dy}{dx}=\frac{x^4+x^2+2}{(x^2+1)^2}$$

Ans. (a) $\dfrac{x^4+x^2+2}{(x^2+1)^2}$

© The Institute of Chartered Accountants of India

174. $\sqrt{\frac{y}{x}} + \sqrt{\frac{x}{y}} = 6$

$\frac{y+x}{\sqrt{xy}} = 6 \Rightarrow x+y = 6\sqrt{xy}$

$\therefore 1+\frac{dy}{dx} = \frac{6}{2\sqrt{xy}}\left(x\frac{dy}{dx}+y.1\right)$

$\left(1-3\sqrt{\frac{x}{y}}\right)\frac{dy}{dx} = 3\sqrt{\frac{y}{x}}-1$

$\frac{dy}{dx} = \frac{3\sqrt{\frac{y}{x}}-1}{1-3\sqrt{\frac{x}{y}}} \Rightarrow \frac{\left(3\sqrt{y}-\sqrt{x}\right)\sqrt{y}}{\left(\sqrt{y}-3\sqrt{x}\right)\sqrt{x}}$

$\frac{dy}{dx} = \frac{3y-\sqrt{xy}}{\sqrt{xy}-3x} = \frac{3y-\left(\frac{x+y}{6}\right)}{\left(\frac{x+y}{6}\right)-3x}$

$\frac{dy}{dx} = \frac{17y-x}{y-17x} = \frac{x-17y}{17x-y}$

Ans. (c) $\frac{x-17y}{17x-y}$

175. ${}^{47}C_4 + \sum_{i=0}^{3} 50-i_{C_3}$

$\Rightarrow \quad 47_{C_4} + 50_{C_3} + 49_{C_3} + 48_{C_3} + 47_{C_3}$

$\Rightarrow \quad 178365 + 19600 + 18424 + 17296 + 16215$

$\Rightarrow \quad 249900$

Ans. (a) 249900

176. $a = 100$

$S_6 = 5.\ S_6$

$\frac{6}{2}[2a+(6-1)d] = 5.\frac{6}{2}[2(a+6d)+(6-1)d]$

$3\ [200 + 5d] = 15\ [200+12d + 5d]$

© The Institute of Chartered Accountants of India

$200+5d = 1000 + 85d$

$80d = -800 \Rightarrow d = -10$

Ans. (a) – 10

177. $S_n = 3n^2 + n$

$\therefore S_1 = 4 \qquad \therefore a_1 = 4$

$S_2 = 14 \qquad a_2 = 14 - 4 = 10 \qquad \therefore d = 6$

$S_3 = 30 \qquad a_3 = 30 - 14 = 16$

$\therefore Tp = a + (p - 1)\, d$

$= \quad 4 + (p - 1)\, 6$

$Tp = (6p - 2)$

Ans. (b) $(6p - 2)$

178. $S_m = S_n$

$$\frac{m}{2}[2a + (m-1)d] = \frac{n}{2}[2a + (n-1)d]$$

$$2ma - 2na = (n^2 - n)d - (m^2 - m)d$$

$2a\,(m - n) = (n^2 - n - m^2 - m)d$

$$2a\,(m - n) = -(m - n)\,(m+n-1)d$$

$\therefore 2a = -(m+n-1)d$

$$S_{m+n} = \frac{m+n}{2}\,[2a + (m+n-1)d]$$

$$\frac{m+n}{2}[(-m+n-1)d + (m+n-1)d]$$

$S_{m+n} = 0$

Ans. (a) 0

179. $-\frac{9}{4}, -2, -\frac{7}{4} \ldots\ldots 0$

$$a = -\frac{9}{4} \quad d = -2 + \frac{9}{4} = \frac{1}{4}$$

$$0 = -\frac{9}{4} + (n-1)\frac{1}{4} \Rightarrow \frac{9}{4} = (n-1)\frac{1}{4}$$

$\Rightarrow 9 = n - 1 \Rightarrow n = 10$

Ans. (b) 10th term

© The Institute of Chartered Accountants of India

180. $6.T_6 = 15.T_{15}$

$\therefore$ 6 (a+5d) = 15 (a+14d)

2a + 10d = 5a + 70d

3a = –60d $\quad \therefore$ a = –20d

T_{21} = a + 20d

= –20 d + 20d

T_{21} = 0

Ans. (c) 0

181. The average of n numbers is x.

If any no.is multiplied to each of datas. Then average will also multiplied by such no.

$\therefore$ New average = (n + 1) x

Ans. (c) (n + 1) x.

182. $Av = \frac{\sum x}{n} \Rightarrow 1.5 = \frac{\sum x}{\delta}$

$= \sum x$ = 12 kg. (increased weight)

$\therefore$ Weight of New person = 65 + 12 = 77 kg.

Ans. (c) 77 Kg.

183. If passes students = x

$\therefore 35 = \frac{39x + 15(120 - n)}{120}$

4200 = 39x + 1800 – 15x

2400 = 24x

$\therefore$ x = 100

Passed Students = 100

Ans. (a) 100

184. $\frac{\sum x}{17} = 45$

$\sum x$ = 765

Total of first 9 numbers = 9 × 51 = 459

Total of last 9 numbers = 9 × 36 = 324

$\therefore$ Value of 9th number = (459 + 324) – 765

© The Institute of Chartered Accountants of India

= 18

Ans. (c) 18

185. $\sum x = 11 \times 30 = 330$

Total of first five numbers = $5 \times 25 = 125$

Total of last five numbers = $5 \times 28 = 140$

$\therefore$ Value of 6th number = (125 + 140) ~ 330 = 65

Ans. (b) 65

186. Ans. (a) Refer properties

187. Ans. (b) Refer Properties.

188. Let A be the hearts playing cards in a part. Let B be the club playing cards in a part.

$\therefore$ $P(A) = \frac{13}{52}$

$P(B) = \frac{13}{52}$

Here A and B are mutually exclusive.

$\therefore$ $P(A \cup B) = P(A) + P(B) = \frac{13}{52} + \frac{13}{52} = \frac{26}{52} = \frac{1}{2}$

189. Total number of balls in the bag = 6 + 4 = 10

Since three balls are drawn out of 10 balls in ${}^{10}C_3$ ways

$\therefore$ Exhaustive number of Cases = $10\,C_3 = 120$

The number of favourable cases two balls are blue and balls is red

= $6C_2 \times 4C_1$

= 60

$\therefore$ Probability of 2 balls are blue and 1 is red = $\frac{6c_2 \times 4c_1}{10c_3}$

= $\frac{60}{120} = ½$

Ans. (c)

190. There are 366 days in a leap year.

Now 366 = 7 x 52 + 2

$\therefore$ The leap year will contain at least 52 Mondays. The possible combination for the remaining two days are:

© The Institute of Chartered Accountants of India

(i) Sunday and Monday

(ii) Monday and Tuesday

(iii) Tuesday and Wednesday

(iv) Wednesday and Thursday

(v) Thursday and Friday

(vi) Friday and Saturday

(vii) Saturday and Sunday

Let A be the event of getting 53 Mondays in the leap year. Therefore, only those combinations will be favourable to the event A which contain "Monday"

$\therefore$ The combination (i) and (ii) are favourable to the happening of A

$\therefore$ P(A) = 2/7

Ans. (a)

191. Given P(A) = 1/3

P(B) = 3/4

and A and B are independent events

$$P(A \cup B) = 1 - P(A \cap B)^1$$

$$= 1 - [P(A^1).P(B^1)]$$

$$= 1 - \{[1 - P(A)] \ [1 - P(B)]\} \quad \text{[A and B are independent]}$$

$$= 1 - \left[\left(1 - \frac{1}{3}\right)\left(1 - \frac{3}{4}\right)\right]$$

$$= 1 - \left\{\left(\frac{2}{3}\right)\left(\frac{1}{4}\right)\right\}$$

$$P \ P(A \cup B) = 1 - \left[\frac{1}{6}\right] = \frac{5}{6}$$

Ans. (b)

192. Out of given 4 letters, there are two letters are vowel (O,E). Let A be the first letter is vowel.

P(A) = 2/4

Let B be the second letter is vowel

P(B) = 1/3

Here A and B are independent

© The Institute of Chartered Accountants of India

P(AB) = P(A).P(B)

$= \frac{2}{4} \cdot \frac{1}{3} = 1/6$

Ans. (a)

193. Out of given 4 letters, there are two letters are vowel (O, E)

Let A be the first letter is vowel.

i.e. P(A) = 2/4

Let B be the second letters is vowel.

P(B) = 1/3

Here A and B are Mutually executive

$\therefore \quad P(A \cup B) = P(A) + P(B)$

$= \frac{2}{4} + \frac{1}{3} = \frac{6+4}{12} = \frac{10}{12} = \frac{5}{6}$

Ans. (a)

194. Let A be the first letter selected M from the 'HOME'.

B be the second letter selected M from the 'HOME'

P(A) = 1/4, P(B) = 1/4

A and B are Mutually exclusive

$\therefore \quad P(AUB) = P(A) + P(B)$

$= \frac{1}{4} + \frac{1}{4} = \frac{2}{4} = \frac{1}{2}$

Ans. (b)

195. By addition thereon

P(A or B) = P(A) + P(B)

0.65 = [1 – P('not A)] + p

= [1 – 0.65] + p

$\therefore$ p = 0.65 – 0.35

p = 0.30

Ans. (c)

197. Since f(x) is a Polynomial.

& a_1, a_2, a_3 are in AP

$\therefore \quad f(a_1), f(a_2), f(a_3)$ also in AP

© The Institute of Chartered Accountants of India

$\therefore$ $f'(a_1), f'(a_2), f'(a_3)$ also in AP

Ans. (a) AP

198. $A = P\left[\frac{(1+i)^n}{i} - 1\right]$

$20{,}000 = P\left[\frac{(1.04)^{10}}{0.04} - 1\right]$

After solving we get

P = 2470 (Approx)

Ans. (a) 2470

199. byx = 1.2 & bxy = – 0.5

This is wrong because bxy and byx have same sign.

Ans. (b) false.

200. The mean of poison distribution is 1.6 and variance is 2. This is wrong because P – d will greater than 2

Ans. (b) false.

© The Institute of Chartered Accountants of India